

UNAIDS/WHO Working Group  
on Global HIV/AIDS and STI Surveillance

# Guidelines on surveillance among populations most at risk for HIV



WHO Library Cataloguing-in-Publication Data

Procurement process resource guide.

(WHO Medical device technical series)

1.Appropriate technology. 2.Equipment and supplies - supply and distribution. 3.Technology assessment, Biomedical I.World Health Organization.

ISBN 978 92 4 150137 8

(NLM classification: WX 147)

**© World Health Organization 2011**

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: [bookorders@who.int](mailto:bookorders@who.int)). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press, at the above address (fax: +41 22 791 4806; e-mail: [permissions@who.int](mailto:permissions@who.int)).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

UNAIDS/WHO Working Group  
on Global HIV/AIDS and STI Surveillance

# Guidelines on surveillance among populations most at risk for HIV

# Acknowledgements

Global surveillance of HIV and sexually transmitted infections is a joint effort of the World Health Organization (WHO) and the Joint United Nations Programme on HIV/AIDS (UNAIDS). The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, provides technical guidance on conducting HIV and STI surveillance. Its mandate is to improve the quality of data available for informed decision-making and planning at the national, regional and global levels.

This document is based on experience in countries and expert reviews from WHO/UNAIDS and the Centers for Disease Control and Prevention (CDC), USA, which have contributed to the writing and revision of the final document. All the persons involved declared no conflicts of interest.

WHO and UNAIDS would like to thank the individuals who contributed to this document including: Donna Stroup, Carolyn Smith, members of the UNAIDS/WHO Working Group on HIV and STI Surveillance, and members of the US Government's Surveillance and Surveys Technical Working Group.

# Contents

Acknowledgements	ii
1 Introduction	3
1.1. Purpose	3
1.2. Background	4
1.3. Classification of HIV epidemics	4
1.4. Ethical issues with populations most at risk for HIV	5
Protect participants' privacy and rights	5
Provide benefits to the community	5
Handle incentives for participation ethically	5
Protect young people	5
Provide protection to participants with illegal behaviours	6
1.5. Process	6
2 Plan surveillance of populations most at risk for HIV	7
2.1. Step 1: Prepare to set up the surveillance system	7
2.1.1 Collaborate with stakeholders	7
2.1.2 Conduct a pre-surveillance assessment	8
2.1.3 Identify populations at increased HIV risk from high-risk behaviour	9
2.1.4 Estimate the size of populations at increased risk for HIV	10
2.1.5 Decide on the geographical areas	10
2.1.6 Consider special issues related to each population of interest	10
2.1.7 Integrate with existing surveillance and monitoring plans	13
2.2 Step 2: Decide on the surveillance design	13
2.2.1 HIV case reporting	13
2.2.2 Programme data used for surveillance	14
2.2.3 Sentinel surveillance	14
2.3. Step 3: Decide on a surveillance sampling strategy	15
2.3.1 Simple random sampling	15
2.3.2 Stratified sampling	16
2.3.3 Cluster or targeted sampling	17
2.3.4 Snowball sampling	17
2.3.5 Respondent-driven sampling	18
2.3.6 Convenience sampling	20
2.3.7 Time–location sampling	20
2.3.8 Sample size	21
2.3.9 Sampling strategies for populations most at risk for HIV	22
3 Conduct surveillance of populations most at risk for HIV	25
3.1 Step 4: Address operational issues	25
3.1.1 Plan staffing and training	25
3.1.2 Develop data collection instruments	26
3.1.3 Plan handling of specimens and HIV testing	27
3.2. Step 5: Data management	27
3.2.1 Plan the data management system	27
3.2.2 Use data management best practices	28
3.2.3 Create a dataset for analysis	28
3.2.4 Document the steps used to create the database	28
3.2.5 Maintain confidentiality and security	29

3.3. Step 6: Data analysis	29
3.3.1 Handling non-response	29
3.3.2 Using weighting	29
3.3.3 Determining HIV prevalence	30
3.3.4 Determining HIV prevalence trends	30
3.3.5 Determining HIV incidence	30
3.3.6 Dealing with uncertainty	31
3.3.7. Ensuring validity	31
4 Use the results/Evaluate the surveillance plan	32
4.1.1 Share the results	32
4.1.2 Develop a dissemination plan	32
4.1.3 Protect the population from stigma or legal action	32
4.1.4 Use the results to improve programmes	33
4.2. Step 8: Evaluate the system	34
Appendix A: List of additional surveillance resources	35
Appendix B: Glossary and acronyms	36
Appendix C: Links	39
References	40

# 1. Introduction

The overall goal of this document is to provide guidance on how to develop and maintain HIV surveillance among populations most at risk for HIV. Ultimately, these surveillance activities should improve the overall understanding of HIV in countries and improve the response to HIV.

This guide complements the second generation surveillance guidelines on how to conduct HIV surveillance activities in low- and middle-income countries. Those guidelines recommend that all countries conduct HIV surveillance among populations with behaviours that increase their risk for HIV, or populations most at risk for HIV infection.

---

## 1.1. Purpose

This document provides guidance on methods for conducting surveillance among populations most at risk for HIV.

Public health surveillance for HIV is the systematic, ongoing collection of data on the occurrence, distribution and trends in HIV infection. In general, the objectives of surveillance include:

- to estimate the magnitude of a health problem in a population at risk
- to understand the natural history of a disease
- to evaluate prevention and control activities
- to monitor changes in trends in the epidemic
- to detect changes in health practices or risk factors
- to identify research needs and facilitate research
- to contribute to the planning process (1).

Additional reasons to conduct HIV surveillance among populations most at risk for HIV are:

- to guide HIV prevention programming at the local level
- to inform priority-setting and resource allocation at the national level
- to contribute to the scientific understanding of HIV transmission in populations most at risk for HIV infection as a result of high-risk behaviour
- to inform disease burden and treatment needs among most-at-risk populations.

When monitoring the HIV epidemic, it is important to identify populations that are most at risk for infection. The sexual and drug-use behaviours of populations contribute to the overall burden of HIV in the country. People with these behaviours are often the first to become infected and are at risk of being infected at a higher rate than those in the general population.

The primary target audience of this guide includes surveillance specialists. In addition, this guide will be useful for programme managers at national and subnational levels to better understand the strengths and weaknesses of the data they are using to make decisions. Finally, this guide will also be useful for donor agencies that support HIV surveillance activities and measure the success of their activities through surveillance data.

Most countries and their development partners already have some form of HIV surveillance among most-at-risk populations. This guide is intended to help refine and standardize their HIV strategies and activities for HIV surveillance among these populations. It presents different approaches along with their advantages and disadvantages. It is hoped that this guide will empower managers and decision-makers to consider additional options for strengthening their HIV surveillance systems so that they are able to compare trends over time.

This guide does not cover all the issues related to HIV surveillance; rather, it serves as a general hands-on reference specifically for surveillance among most-at-risk populations. It cites additional materials and resources for further information on surveillance and includes country examples.

## 1.2. Background

This document complements a number of other guidelines that provide further information on specific surveillance activities. A complete list of reference materials is provided in Appendix A. These guidelines are available on the UNAIDS and WHO web sites.

Some of the core sources include the following:

- *Guidelines for estimating the size of populations most at risk to HIV* presents a process for creating local and national size estimates.
- An update to *Guidelines on second generation surveillance* explores new tools and techniques from a field perspective to enhance the first guideline published in 2000.

There are a number of documents that describe monitoring and evaluation activities among populations most at risk for HIV, including the *Framework for monitoring and evaluating HIV prevention programmes among most-at-risk populations* (2007). ([http://data.unaids.org/pub/Manual/2008/jc1519\\_framework\\_for\\_me\\_en.pdf](http://data.unaids.org/pub/Manual/2008/jc1519_framework_for_me_en.pdf)). Operational guidelines for the monitoring and evaluation of HIV-related programmes targeting persons who inject drugs and men who have sex with men are anticipated in 2011.

For more in-depth information on the topics covered here as well as a self-teaching manual on surveillance among most-at-risk populations, see *Surveillance of most-at-risk populations* ([http://globalhealthsciences.ucsf.edu/PPHG/surveillance/surv\\_modules.html](http://globalhealthsciences.ucsf.edu/PPHG/surveillance/surv_modules.html)).

## 1.3. Classification of HIV epidemics

UNAIDS/WHO and partners have identified three epidemic categories to help countries focus surveillance activities. These categories are suggestions. Countries should know best where new infections are coming from and what type of epidemic exists in their country.

Each country has a unique epidemic and usually has multiple subepidemics within different parts of the country. Because of the diversity among HIV epidemics, it is critical to “know your epidemic”. That means understanding how the epidemic differs within subpopulations and in geographical areas. Surveillance data will provide the information to allow programme managers to better know their epidemics. Moreover, they will allow programme managers to respond more effectively to the epidemic. As more surveillance data become available, surveillance officers should evaluate their surveillance system to ensure that it is appropriate for the type of epidemic. Table 1.1 will help determine where to focus the surveillance system and whether the focus needs to shift over time.

**Table 1.1. Base surveillance activities on the type of epidemic**

Epidemic state, situation	Surveillance focus
Low-level ■ HIV has not reached significant levels in populations most at risk for HIV infection as a result of high-risk behaviour. ■ HIV is largely confined to people within populations most at risk for HIV infection as a result of high-risk behaviour.	■ Focus surveillance activities in populations most at risk for HIV.
Concentrated ■ HIV has spread rapidly in one or more populations most at risk for HIV infection as a result of high-risk behaviour. ■ The epidemic is not yet well established in the general population.	■ Continue surveillance among most-at-risk populations. ■ Begin surveillance activities in the general population, especially in urban areas.

<p>Generalized</p> <ul style="list-style-type: none"> <li>■ The epidemic has matured to a level where transmission occurs in the general population, independent of populations most at risk for HIV.</li> <li>■ Without effective prevention, HIV transmission continues at high rates in populations most at risk.</li> <li>■ With effective prevention, prevalence will drop in populations most at risk before they drop in general population.</li> </ul>	<ul style="list-style-type: none"> <li>■ Focus routine surveillance on the general population.</li> <li>■ Conduct surveillance among populations most at risk for HIV.</li> </ul>
--	---

As described in Table 1.1, each type of epidemic should maintain some surveillance of populations most at risk for HIV. Use these guidelines to facilitate surveillance in all settings.

#### 1.4. Ethical issues with populations most at risk for HIV

Collection of any data requires attention to human subjects' concerns. Before considering surveillance of populations most at risk for HIV, the following principles should be considered.

##### **Protect participants' privacy and rights**

In many cases, populations that are the target of HIV surveillance are vulnerable and might be accorded special protection under ethical regulations. Handling of data on risk behaviours and HIV status may increase the risk of harm to these populations due to stigma, economic loss or legal liability (2).

Participating in surveillance activities should either be voluntary or, in specific circumstances, should be based on unlinked anonymous testing of blood samples. Unlinked anonymous testing should be used as a surveillance strategy only when data from clinical settings and other studies cannot provide an accurate measure of the prevalence or trends of HIV infection (3). Respect each person's right to privacy. It is critical to offer privacy and assurances of confidentiality during data collection. Keep data confidential. In turn, the perception of confidentiality will influence the completeness of reporting and disclosure of risk.

##### **Provide benefits to the community**

Public health research should always provide a benefit to society. Extend the ethical principle of *do no harm* to providing benefits to surveillance subjects. Whenever possible, provide individuals participating in the survey with:

- test results
- information about HIV and AIDS
- counselling (on preventing HIV or on other health or social needs)
- treatment and care to the extent possible with local resources
- referrals to other services.

Consider how the results will be used before starting a surveillance activity among most-at-risk populations. The results should be used for the good of public health while minimizing any harm to the participants.

##### **Handle incentives for participation ethically**

Providing benefits in the form of cash payment or vouchers for goods and services is often a useful recruitment and retention technique (4). However, the use of incentives should be considered carefully during the ethical review. Any incentive must consider the customs of the community but should not be so large that it is considered coercive or exploitative (5). The ethical concern here is that the offer of financial compensation, medical services or some goods might cause participants to consider only their short-term benefits and they might underestimate the long-term costs. This might entice some participants to provide data and could create a bias to the results (6).

##### **Protect young people**

When surveillance of most-at-risk populations includes collecting data from adolescents, additional ethical issues arise. Young people may be particularly vulnerable to exploitation, abuse and other harmful outcomes (legal, physical and social). Follow the guidelines for collecting data on adolescents (4,7).

### Provide protection to participants with illegal behaviours

Safeguard participants from situations in which they might face legal persecution. This is critical for HIV surveillance in any context. Some of the behaviours that cause increased risk for HIV are illegal in some settings. Legal protection of confidentiality are constantly in flux (8). This can be avoided by including members of the population who can provide information on the local context in surveillance pre-planning.


For additional information, see *Ethical issues to be considered in second generation surveillance* ([http://www.who.int/hiv/pub/epidemiology/en/sgs\\_ethical.pdf](http://www.who.int/hiv/pub/epidemiology/en/sgs_ethical.pdf)). Additional guidelines on ethics and HIV surveillance will be available on the WHO web site shortly.

---

## 1.5. Process

Planning and conducting surveillance among populations most at risk for HIV can be summarized by the eight steps shown below.

**Figure 1.1. Planning and conducting surveillance among populations most at risk for HIV**




The remainder of this document discusses the processes shown in Figure 1.1. Guidance and examples are provided on methods:

- to plan surveillance activities, steps 1–3, Section 2
- to conduct surveillance, steps 4–6, Section 3
- to disseminate results and evaluate the surveillance system, steps 7–8, Section 4.

## 2. Plan surveillance of populations most at risk for HIV

The first three steps of the process are general preparations for conducting surveillance among populations most at risk for HIV. Some countries might have an existing surveillance system. It might still be useful to review the first steps to ensure that the system is well designed.

**Figure 2.1. Plan HIV surveillance among populations most at risk for HIV**



### 2.1. Step 1: Prepare to set up the surveillance system

While preparing the surveillance system, it is important to develop a protocol to ensure the success of the surveillance system. This will help ensure that accurate, appropriate and useful data will be captured by the surveillance system. Be sure the protocol covers the following areas:

- how surveillance stakeholders will be involved in activities
- planned pre-surveillance assessment
- populations of interest, with explicit definitions of these populations, based on the behaviours that put them at increased risk for HIV
- the size of the population or populations
- geographical boundaries, sites or location of surveillance
- coverage and descriptions of the representativeness of the population
- frequency of surveillance
- how surveillance activities will be merged with other monitoring and evaluation plans.

#### 2.1.1 Collaborate with stakeholders

HIV surveillance cannot be done in isolation. Use a team approach to improve the quality and credibility of the results. Table 2.1 provides a list of possible stakeholders and what they can bring to the surveillance activities. Plan to involve members of HIV prevention programmes, monitoring and evaluation programmes, and resource planning structures. Identify a team of stakeholders to help in designing and implementing HIV surveillance activities.

**Table 2.1. Potential stakeholders to include in planning surveillance activities among populations most at risk for HIV**

Stakeholder	What they bring
Members of the populations most at risk for HIV	<ul style="list-style-type: none"> <li>■ Provide insight into: <ul style="list-style-type: none"> <li>• the timing of surveillance</li> <li>• the appropriateness of the sampling strategy</li> <li>• the wording for the questionnaire, and</li> <li>• instrument design</li> </ul> </li> <li>■ Provide study legitimacy in the community</li> <li>■ Provide the legal and social context</li> </ul>
Nongovernmental organizations or civil society organizations working with most-at-risk populations	<ul style="list-style-type: none"> <li>■ Provide information on the needs of most-at-risk populations</li> <li>■ Have experience in reaching the population to inform the sampling strategy and instrument design</li> </ul>
HIV prevention experts	Ensure that the surveillance results provide information on a timely basis for programme planning or advocacy
Survey and census implementers	<ul style="list-style-type: none"> <li>■ Advice on sampling strategies</li> <li>■ Additional data for size estimation activities</li> </ul>
Monitoring and evaluation staff	<ul style="list-style-type: none"> <li>■ Ensure that surveillance activities are part of the national monitoring and evaluation plan</li> <li>■ Reduce the chances that efforts will be duplicated by different organizations</li> <li>■ Combine data/analysis of results with other monitoring and evaluation activities</li> <li>■ Disseminate combined results</li> </ul>
Staff engaged in surveillance for sexually transmitted infections	Improve efficiency by combining HIV surveillance activities with surveillance for sexually transmitted infections

### 2.1.2 Conduct a pre-surveillance assessment

Conduct a pre-surveillance assessment before starting surveillance activities. The purpose of this assessment is to understand the epidemic in the country. Review existing HIV and STI surveillance results as well as findings from behavioural surveys. Consider which populations are at increased risk for HIV and how their behaviours put them at increased risk. Guidelines on how to conduct a pre-surveillance assessment are available at: [http://data.unaids.org/pub/Manual/2005/20050101\\_gs\\_guidepresurveillanceassmnt\\_en.pdf](http://data.unaids.org/pub/Manual/2005/20050101_gs_guidepresurveillanceassmnt_en.pdf).

In brief, pre-surveillance activities should gather information on the subpopulations and geographical areas to be included in the surveillance. The assessment should include qualitative fieldwork

- to identify and verify hotspots,
- to verify the definition of the population, and
- to guide the planning and logistics of surveillance fieldwork.

The pre-surveillance assessment will ensure that the right subpopulations are being considered and are defined correctly. If the population is defined incorrectly, the surveillance team might fail to detect emerging epidemics and miss the opportunity to target interventions where they will make the most difference.

Qualitative research is necessary to describe the patterns of behaviour, points of access and barriers to surveillance for the populations (9). Surveillance methods will differ depending on whether the person is self-identified and, if so, whether the person is part of a subculture (10).

It is worth taking the extra time and effort to go through the pre-surveillance process to plan appropriately and refine the surveillance protocol. Conducting the assessment could help avoid mistakes that might ultimately cost lives, time and money.

### 2.1.3 Identify populations at increased HIV risk from high-risk behaviour

Specific behaviours put individuals at increased risk for HIV. During the pre-surveillance assessment, the available resources are compiled to help identify populations with these behaviours. Very high transmission rates are found in groups that engage in the following behaviours:

- inject drugs with used needles
- have anal sex
- have sex with many partners without protection.

Specific populations with behaviours that put individuals at increased risk for HIV have been identified as being important for HIV surveillance (11):

- Sex workers
- Clients of sex workers
- People who inject drugs
- Men who have sex with men.

Identify the actual behaviour that puts people at increased risk for HIV when selecting subpopulations for surveillance activities. For example:

- Sex workers have large numbers of partners, often within a very short time, increasing their exposure to HIV and sexually transmitted infections. Sex workers are often at a disadvantage in negotiating condom use because their partners pay for their services.
- Clients of sex workers are also at increased risk for infection because of the sex workers' high turnover of sex partners. Clients of sex workers often act as a bridge to low-risk populations, which means that they are often the link between the populations with behaviours that put them at increased risk for HIV and low-risk populations. For example, a client of a sex worker may transmit HIV to his regular partner.
- Persons who inject drugs are at increased risk because of the high transmissibility of HIV when injecting equipment is shared. When a needle, syringe or other injecting equipment is shared, the blood from the first injector is often still in the equipment. It is injected into the next user's body. This is a very efficient method for transmitting HIV or hepatitis B and C.
- Men who have sex with men are often identified as another population at increased risk for HIV. This is because unprotected anal sex has much higher transmissibility than vaginal sex.
- Women who receive anal sex are also at increased risk for HIV.

Carefully consider the difference between vulnerable populations (such as persons with disabilities) and populations with behaviours that put them at increased risk. Conducting surveillance among vulnerable populations whose risk-taking behaviours are not well understood is not useful (although it might be useful for programming for those populations).

Transgender persons often have unique networks of sexual partners or clients. The risk behaviours are often the same as described above – anal sex or selling sex – but they take place in select venues or with select populations. Given the high levels of HIV prevalence among this population, it is important to carry out special surveys to estimate the size of this population and estimate the HIV prevalence in this population.

Transgender populations can vary from an individual who dresses as the other sex (transvestite) to persons who have had surgery to modify body parts (trans-sexual). The population is difficult to categorize and often identifies with different terminology. Formative research is critical among this population to ensure that the correct terms and questions are included in any survey. Participation from the community is also essential. For guidance on how to conduct surveys among transgender populations, see <http://transpulse.ca/documents/OHTN-%20Best%20Practices%20for%20Trans%20Participants-%20GBauer%202009%20vFINAL.pdf>

Decide on a clear and specific definition of the populations that will be covered with surveillance activities. Here are some examples:

- Sex workers – “Women who have received money for sex at a brothel in the past six months”
- Persons who inject drugs – “Persons who have injected any non-medical drug in the past three weeks”.

Definitions will depend on the surveillance strategy and design so no global guidance can be provided on these definitions. Keep in mind that definitions should be precise, time-bound, measurable, valid and reliable.

### **Proxy populations**

It is often necessary to use proxy definitions for at-risk populations that are not a distinct social group. A *proxy definition* uses a sociodemographic characteristic of a group, such as occupation, or places where persons engaging in risk behaviours are likely to be found (such as men at bars, male migrants living in dormitories, etc.).

The proxy definition is not the cause of the increased risk for HIV. For example, *truck drivers* are often used as a proxy definition for clients of sex workers because some studies show that a higher proportion of truckers are reported to be clients of sex workers than men in the general population. However, on its own, driving a truck is not a risk for acquiring HIV. Similarly, *prisoners* are often a proxy definition for most-at-risk populations. In some settings, prisoners have been found to inject drugs or have anal sex.

A proxy definition is almost always imperfect:

- Some people who meet the proxy definition may not engage in the risk behaviour.
- Some people who have the risk behaviour may not meet the proxy definition.

A proxy definition is useful only if there is evidence that a large proportion of individuals in the group practise the high-risk behaviour of interest. When using data from proxy groups to describe the epidemic, be clear why a proxy group has been adopted. Document any local data which demonstrate that the proxy group does define a population with higher risk behaviours.

### **2.1.4 Estimate the size of populations at increased risk for HIV**

Estimates of the size of at-risk populations are critical for interpreting surveillance results and can be used to justify surveillance activities to stakeholders and funders. Estimates can be a strong force in policy change and resource allocation by describing the potential population at risk. In addition, estimates of population size inform the development, implementation, and monitoring and evaluation of interventions.

Population sizes can often be estimated at the same time as surveillance activities. Information on how to conduct size estimation studies is available in the *Guidelines on estimating the size of populations most at risk to HIV (12)*.

### **2.1.5 Decide on the geographical areas**

Population members in different regions of the country may have different behaviours and thus different risks. These differences are important for planning surveillance efforts.

Differences by region may make it difficult to generalize surveillance results from one city to the rest of the country. Surveillance activities will be needed in multiple locations within the country to ensure that these differences are captured.

Again, the information gathered during the pre-surveillance assessment should assist in deciding on the geographical regions where surveillance will be conducted.

### **2.1.6 Consider special issues related to each population of interest**

#### **Sex workers**

Sex workers – people who exchange sex for money or goods – are an important group in the HIV epidemics of many countries, especially in countries where condom use by sex workers and their clients is low. Sex workers may have a higher prevalence of HIV than other populations because of their high turnover of sexual partners and financial incentives by clients to not use condoms.

Sex workers differ markedly from one place to another and by the level of the HIV epidemic (13). In addition, there may be many different categories of sex work. In pre-surveillance activities, it is especially important to get the community to support the research so that they can provide input on the surveillance design.

In addition, researchers should forge alliances with individuals and organizations in a position to deliver specific intervention programmes. These include:

- sex workers' groups
- local authorities
- gatekeepers, such as brothel owners
- women's groups.

Data that should be collected in addition to HIV status include: frequency of sex work, time since initiation in sex work, frequency of condom use, non-client partners, questions on the reach of prevention and care programmes available in the area, other high-risk behaviours and other relevant variables.

### ***Clients of sex workers***

Clients of sex workers are at increased risk of infection because they have sex with sex workers who have large numbers of sex partners. As described earlier, clients of sex workers act as a bridging population to the general population. Surveillance among clients of sex workers is increasingly being conducted among clients recruited at sex work venues (14,15, 16).

Surveillance of client populations can also be done using proxy populations that are believed to have sexual contact with sex workers. These include:

- commercial drivers
- military personnel
- patients at STI clinics who report visiting sex workers.

In addition to HIV status, some additional data which should be collected among clients include: frequency of visits to sex workers, sexual behaviour with regular partners, condom use frequency, other high-risk behaviours and other relevant variables.

### ***Persons who inject drugs***

In most societies, the illegal nature of injecting drug use and its associated stigma mean that persons who inject drugs are often a hidden population (17). Implementing surveillance among persons who inject drugs requires some general principles:

- Combine HIV surveillance activities with a larger surveillance framework, including other health issues such as hepatitis B and C surveillance.
- Work with organizations that have links to drug users:
  - former users
  - drug-abuse treatment programmes
  - law enforcement settings
  - community outreach workers who often are recruited from among current users
  - treatment and rehabilitation centres (18).

In many countries, the population of those who inject drugs varies from year to year, so plan to conduct size estimation studies regularly rather than relying on old data. As described earlier, preventive services for persons who inject drugs should accompany HIV surveillance activities (19).

Collect data on risk and preventive behaviours, including information on screening and vaccination for hepatitis. Include needle-sharing, sexual activity (such as condom use), access to free sterile injection equipment, participation in prevention activities and access to antiretroviral treatment (20, 21). See *Technical guide for countries to set targets for universal access to HIV prevention, treatment and care for injecting drug users*. ([http://www.who.int/hiv/pub/idu/idu\\_target\\_setting\\_guide.pdf](http://www.who.int/hiv/pub/idu/idu_target_setting_guide.pdf))

Different drug-injecting behaviours may contribute to HIV transmission, for example:

- needle-sharing
- indirect sharing ("backloading" or "frontloading")
- sharing filters or cookers.

In *indirect sharing*, one syringe is used to prepare the drug solution, which is then divided into one or more syringes for injection. The solution is transferred between syringes with the needle removed (*frontloading*) or plunger removed (*backloading*). This allows virus transmission if the preparation syringe is not clean.

The importance of each of these methods of sharing may depend on local behaviour. For example, if direct sharing has been virtually eliminated due to intervention programmes and behaviour change, then surveillance should monitor indirect sharing.

Surveillance should also monitor sharing that involves many different “sharing partners” within a short period (“rapid partner-change” behaviour), since this behaviour is associated with rapid transmission of HIV among persons who inject drugs. Technical guidelines that provide specific guidance on monitoring and evaluating interventions for injecting drug use are available. ([http://www.who.int/hiv/pub/idu/idu\\_target\\_setting\\_guide.pdf](http://www.who.int/hiv/pub/idu/idu_target_setting_guide.pdf))

Finally, collect information on access to sterile injecting equipment. Easy access to sterile injection equipment is critical for HIV prevention among persons who inject drugs.

In summary, surveillance among persons who inject drugs should incorporate:

- data on drug injection risk behaviour
- sexual risk behaviour
- size and growth of the population
- patterns of drug use
- access to sterile injecting equipment
- participation in prevention activities, including screening for hepatitis
- access to antiretroviral therapy.

### ***Men who have sex with men***

HIV among men who have sex with men is increasing in many countries of the world (22,23,24). Surveillance in this population has often underestimated the magnitude of infection and risk among this population. Many countries do not include men who have sex with men in their regular surveillance activities because homosexuality is illegal and highly stigmatized (25).

The impact of risk among men who have sex with men varies, depending on the level of the epidemic. Even in some regions with a generalized epidemic, men who have sex with men may have a higher HIV prevalence than among the general population (26).

Use formative research to enlist the cooperation of the community.

Behavioural questions that should be included in surveys with men who have sex with men include: the number of partners, sexual behaviour and frequency of sex with female partners, time since initiation of male-to-male sex, frequency of condom use, participation in HIV prevention or care programmes and other relevant variables.

### ***Other populations***

In some countries, access to populations most at risk for HIV may be limited. As a result, surveillance activities may be planned among proxy populations. In some cases, regular and consistent HIV testing and counselling will be available from proxy populations (for example, prisoners), providing an opportunity to analyse potential trends. It is important to ensure that ethical processes were followed to generate these data before using them.

Surveillance systems based on proxy populations might not be adequate. Although they can be useful for supplementing surveillance, they will typically not provide full and representative information on the actual group of interest. As a result, they suffer from the same problems and weaknesses of such sampling or selection biases (sampling biases and weaknesses are described later in this document). Additional variables that should be asked of these populations depend on the behaviour that puts the group at increased risk.

***Incarcerated populations***

Persons in prison might be more likely to engage in behaviours that increase their risk for acquiring HIV through injecting drugs, engaging in anal sex with fellow prisoners (27,28) or tattooing (29). While these behaviours occur in many prison settings, incarcerated populations are not a proxy population for all injecting drug users or men who have sex with men.

Prison populations are often used as proxy populations. Surveillance among prisoners should include the reason for each prisoner's incarceration (this can provide potential information to determine if they were infected before imprisonment).

Keep in mind that incarcerated persons may have been infected before they were incarcerated.

***Military recruits***

In some countries, military recruits are more likely to visit sex workers. This makes them a useful proxy population for clients of sex workers. In addition, this population is often routinely tested for sexually transmitted infection (including HIV) through existing medical systems.

However, being in the military is not on its own a risk factor for HIV infection so interpret results with caution.

**2.1.7 Integrate with existing surveillance and monitoring plans**

After conducting all of these preliminary activities, including the formative assessment and size estimation, revise and review the HIV surveillance protocol. Ensure that the protocol documents the decisions made and plans future surveillance activities on a timeline. Make sure that the surveillance protocol is incorporated into the national monitoring and evaluation plan to avoid overburdening staff and duplicating efforts. Use the surveillance protocol to develop funding requests for surveillance activities.

**2.2 Step 2: Decide on the surveillance design**

Surveillance results will help determine:

- how many people are infected with HIV among the population most at risk, and
- the factors that might be associated with their HIV status.

Therefore, surveillance activities must collect biological and behavioural data.

There are a few ways to collect biological data. Options include:

- HIV case reporting
- using programme data for surveillance
- sentinel surveillance.

Regardless of the biological surveillance design, behavioural data should also be collected. Tracking HIV prevalence or incidence may show the progress in implementing prevention programmes. However, these data alone will not indicate why the programmes succeeded or failed. Behavioural data can show associations with particular aspects of risk.

Historically, surveillance activities for biomarkers and for behavioural indicators were conducted as separate activities. When planning surveillance activities that collect information on biomarkers, it is currently recommended to also collect data on risk and preventive behaviours.

**2.2.1 HIV case reporting**

Many developed countries rely on HIV case reporting (and the self-reported transmission route) for surveillance among most-at-risk populations. However, this method has several limitations:

- Many individuals are not diagnosed until long after they have been infected, or not at all, leading to underreporting.
- The self-reported mode of transmission is potentially biased, as individuals may not want to disclose high-risk behaviours to health-care providers.
- HIV case reporting generally provides only limited data on the risks of HIV transmission or behaviour.

Therefore, if HIV case reporting will be used, it is important to conduct alternative or additional surveillance activities to supplement the case reporting data.

Because of the time lag of almost a decade between infection and development of AIDS, AIDS case reporting does not reflect the recent dynamics of the HIV epidemic. In addition, as the number of people receiving antiretroviral therapy increases, the number of new AIDS cases will decline. AIDS case reporting is therefore not very useful for surveillance. Instead, building on systems for the delivery of antiretroviral therapy, surveillance of advanced HIV is recommended (30).

Case reporting often needs to be based on names or national identity numbers to avoid double counting. If individuals are required to report on behaviours that might be illegal or highly stigmatized, storing identifying information along with self-reports of those data might be unethical.

### **2.2.2 Programme data used for surveillance**

Many countries collect biological results from their programmes, such as sexually transmitted infection (STI) clinics or HIV testing and counselling services. Routine programmes often focus on populations at increased risk for HIV.

STI clinics or voluntary counselling and testing services collect information on risk behaviours. This allows programme managers to separate the test results of men who have sex with men, women who have anal sex, sex workers and persons who inject drugs. Annual measures of the HIV prevalence in these separate populations provide an indication of HIV trends in most-at-risk populations.

However, data from voluntary counselling and testing and other services are often biased, depending on who decides to participate in these services. For example, when the services charge fees, these services will be attended by more wealthy members of the population. Moreover, the bias is likely to change over time as more or less people access these services as programmes expand or undergo budgetary restrictions. Similar to case reporting, this method relies on self-reported behaviours provided in a setting that might not be confidential. Data may be biased depending on the stigma attached to the behaviour in the country.

### **2.2.3 Sentinel surveillance**

Sentinel surveillance uses selected sites for collecting measures of HIV prevalence in a population.

Sentinel surveillance can take the form of:

- a community survey or
- a facility-based survey.

Both require the identification of subgroups of the population who are at increased risk. The section on sampling later in this document describes the methods available for sentinel surveillance.

Similar to case reporting, this method relies on self-reported behaviours to determine the mode of transmission. In many survey settings, interviewers are unknown to the respondent. This ensures some level of confidentiality.

Sentinel surveillance at the community level has several challenges. All of these factors influence the validity of the data and the generalizability of the data to a larger population.

- Sampling of most-at risk-populations is often challenging because no complete *sampling frame* (list of members of the population) is available. Because of this, the resulting sample may be biased and difficult to replicate over time.
- When random selection rules are not used, interviewer and self-selection sampling biases may result. As a result, the survey may under-sample members of a given population who do not occupy obvious niches in the community or who do not approach outreach workers. In other words, the survey may not reach hidden subpopulations.

- Sampling of subgroups of interest may only recruit those who are at higher risk for HIV infection and only later recruit those at lower risk (31).
- In sampling populations of interest, geographical areas may not be sampled in proportion to numbers of the population of interest. For example, drug users may not be sampled in proportion to the intensity of drug use in the region. The probability of selecting drug users may not be known.

All of these issues are discussed further in the section on sampling.

---

### 2.3. Step 3: Decide on a surveillance sampling strategy

Surveillance is conducted to allow comparisons of data from the same group over time. Thus, samples should be:

- representative of the population
- able to be repeated (or replicable) in a later survey.

For most-at-risk populations, it is difficult to replicate surveys. Populations may be difficult to reach (hidden) or transient; that is, people move into or out of the population. To address these issues, HIV prevalence studies among most-at-risk populations often use probability, convenience or snowball sampling methods, which are not always replicable over time.

While selecting a sampling strategy, consider the following:

- It should be feasible to sample from the same population in future surveys.
- The sample should be as representative of the population as possible.
- The sampling strategy should match available resources for this round and future rounds of the survey.
- It should be possible to carry out the survey (reaching all persons in the sample) in a reasonably short period of time.
- The cost involved should be affordable.
- Legal and ethical issues of implementing the sampling strategy should be taken into account.

#### 2.3.1 Simple random sampling

##### **Description**

The way a sample is selected determines the scientific viability of the study. A simple random sample is the gold standard for sampling considerations. Simple random sampling assumes that everyone in the target population has the same, known probability of being included in the sample, like drawing numbers or names from a hat. More commonly, each member of the population is identified by a number and random number tables or computer algorithms are used to select the required sample size.

##### **Advantages**

From a statistical perspective, simple random sampling:

- eliminates bias from a sample of volunteers who might have different characteristics than the complete population
- provides each member of the population with the same chance of being selected
- provides a sample that contains members with characteristics similar to the population as a whole.

##### **Limitations**

As straightforward as this seems, implementing a true simple random sample in fieldwork can be expensive, inconvenient or impossible:

- Implementing a random sample requires a *sampling frame* or list of the target population. Clearly, this requirement is not met for many populations at increased risk for HIV. (See Box 1.)
- Even if a list does exist, the sample population might be dispersed throughout a geographical area, making it expensive and time-consuming to carry out.
- A random sample may miss people or behaviours that are not common in the population.

In cases where a sampling frame and easy access to the population are not available, alternatives are necessary. Excellent reviews of conventional sampling techniques can be found elsewhere (32,33,34). Several variants of classical sampling are important for surveillance among populations at increased risk for HIV (35). These variants are presented below.

#### **Box 1: What is a sampling frame?**

A sampling frame is simply a list of all of the individuals eligible to be in the survey. Persons participating in the survey are selected from the sampling frame and are considered the survey sample. An incomplete sampling frame can result in a biased sample and inaccurate results if the people excluded from the frame have different characteristics than those included in the sampling frame.

### **2.3.2 Stratified sampling**

#### **Description**

Stratified sampling is modified simple random sampling. Stratified sampling may provide an economic saving over simple random sampling:

- Divide the target population into groups (strata) that are more similar within the group than the groups are to each other.
- Now take a simple random sample from each group independently.

Define strata by the nature of the sampling question. For example, if the variable of interest is condom use, and different types of sex workers used condoms at different frequencies, the strata could be defined as the various types of sex workers (street-based, brothel-based, home-based) in the population.

#### **Advantages**

When data are collected in this way, statistical methods allow weighting schemes to be used to estimate the variability and potential error of the survey due to the sampling design (36). (See Box 2.)

#### **Limitations**

It is important to have a clear understanding of the population and the differences within the population to be able to stratify it. In the example above, we would need to know the different types of sex workers and their behaviours. Once the strata are known, a sampling frame is needed for each stratum. Choose the sample from these lists.

#### **Box 2: How to weigh survey results**

When a sample is drawn from a sampling frame, it is common to draw more from one section of the population than another. This is referred to as stratified or cluster sampling.

When stratified sampling is used, it is possible to correct the bias in the sample by calculating a variable that truly represents the population of interest. For example:

- If one subgroup of the population is *overrepresented* in the survey, give this subgroup a weight of less than one.
- If a subgroup is *underrepresented* in the survey, give them a weight greater than one.
- The overall weighted frequency is equal to the total number of people in the survey.

All survey variables are then calculated using these weights to correct for the biases introduced from the sampling strategy. In addition, weights are used to calculate the precision of the estimate and the potential error as a result of the survey design or the design effect.

### 2.3.3 Cluster or targeted sampling

#### **Description**

Cluster or targeted sampling is also a modified simple random sampling. The target population may be naturally divided into groups (as with stratified sampling). Within the groups there is diversity in the variable of interest but, on average, one group is similar to the other. Each cluster should be a fairly good representation of the entire population on a smaller scale. Often, clusters are geographical areas, such as:

- villages
- regions of a rural area
- sections of a town or city
- physical units such as brothels, bars or streets.

To conduct cluster sampling, randomly select the clusters and then perform simple random sampling within the clusters (37).

#### **Advantages**

As with stratified sampling, weighting allows for calculation of the **sampling error** (38). Using the sex worker example, if most of the population of interest is street-based, cluster sampling may be appropriate. Cluster sampling commonly uses maps to guide the stratification. This allows for a greater generalizability to a geographical area.

#### **Limitations**

It is usually easy to break groups into clusters when using geographical areas. Use mapping of the population to be sure that a sample of the entire population in the region is included. A sampling frame from which to draw the sample is needed for each cluster.

As described earlier, stratified and cluster samples can use weighting schemes to create statistical estimates of variability and errors. This means that they can produce **probability samples**. Properly conducted, these adaptations can produce estimates with as much or more accuracy as simple random sampling but with a substantial saving of resources.

However, these designs could lead to higher estimates of error compared with that from simple random sampling. When surveys are structured with stratifications or clusters, they are described as having a **design effect**. The design effect of the study needs to be considered when planning the sampling strategy and calculating sample sizes. Involve a statistician on the surveillance team for these types of issues.

The first three options described above are often not feasible when working with most-at-risk populations. There are usually no sampling frames if the entire most-at-risk population in a geographical area is considered the sampling universe. Therefore, non-probability sampling is often necessary for surveillance of most-at-risk populations.

### 2.3.4 Snowball sampling

#### **Description**

Snowball sampling is one type of non-probability sampling. Persons already interviewed or measured are used to find study subjects. Generally, subjects first contacted are asked to name acquaintances who are then approached, interviewed and asked for additional names. In this way, a sufficient number of subjects can be accumulated to give a study adequate power (39).

For example, if the goal is to study persons who inject drugs, sampling from needle exchange programmes, while accessible, will miss many women, youth and those who have recently started injecting. This would produce a statistically representative sample of an unrepresentative part of the target population. Snowball sampling was developed to address this problem by reaching individuals in diverse social networks.

#### **Advantage**

The method allows the sampling of populations that are networked when there is no sampling frame.

### **Limitations**

Snowball sampling has many limitations.

- If the goal is repeated surveillance on the same group, snowball sampling will not be appropriate. There can be no way to compare the results over time.
- Bias can result since most people tend to name acquaintances who are similar to themselves demographically: by race, ethnicity, education, income or religion.

## **2.3.5 Respondent-driven sampling**

### **Description**

Respondent-driven sampling is a modification of snowball sampling which uses a mathematical model to weight the sample data to compensate for the fact that the sample is not a simple random sample (40,41). Respondent-driven sampling is an important method for sampling hard-to-reach groups, including when the groups may be small relative to the general population or when a list (sampling frame) of population members does not exist.

To conduct respondent-driven sampling:

- Identify and recruit seeds (initial survey respondents). The number of seeds required varies by the total sample size required and the diversity of the target population.
- Give each seed coupons and ask them to recruit three to five acquaintances.
- Each recruit who visits the study site brings a coupon that identifies (by number) who referred them.
- No names are provided, just study identification numbers combined to develop a list of the numbers of acquaintances per study participant.
- Give each recruit coupons and ask them to recruit three to five acquaintances.
- Recruiting may go on for several waves.
- A mathematical model based on the Markov chain theory and network theory weights the sample to address the probability that people will refer their friends, making the sample less random.

### **Advantages**

The resulting statistical theory controls for the biases associated with chain-referral methods, providing both population estimates and estimates of variability for those estimates (42).

Individuals are accessed through their social networks, making it more likely to reach those who shun large public venues and avoid the street.

### **Limitations**

As a method, respondent-driven sampling is costly and time-consuming. Sources of significant bias can exist if the population is not well networked. Entire sections of the population of interest can be missed if they are not connected to the initial seeds.


Depending on the number of initial seeds, the start-up of a respondent-driven sampling study can be slow.

Respondent-driven sampling requires specific software for analysis (available at: <http://www.respondentdrivensampling.org/main.htm>). Recent research has found that there is large variation in the estimates produced through respondent-driven sampling. Large sample sizes should improve on the estimates (43).

There have been mixed results using respondent-driven sampling with different most-at-risk populations. In some contexts, the populations were not well networked, making respondent-driven sampling a difficult sampling strategy.

A more detailed description of how to conduct respondent-driven sampling is available at: <http://www.respondentdrivensampling.org/>.

**Figure 2.2. Respondent-driven sampling recruitment network for one seed by type of female sex worker in Viet Nam**



Source: Johnston LG et al. Effectiveness of respondent-driven sampling to recruit female sex workers in two cities in Vietnam. *Journal of Urban Health*, 2006, 83 (Suppl 7):16–28. <http://www.springerlink.com/content/j31w862u2v65538n>

### Respondent-driven sampling in Beijing, China

Respondent-driven sampling was used for three rounds of surveys to recruit men who have sex with men in Beijing, China in 2004, 2005 and 2006. The surveys included a structured interview to collect behavioural data and blood specimens to test for HIV, syphilis, hepatitis B virus and hepatitis C virus. The number of participants in the surveys was 325, 427 and 540 in the 2004, 2005 and 2006 surveys, respectively. In the 2004 survey, only one seed was recruited; in the 2005 and 2006 surveys, 10 and 8 seeds were recruited, respectively.

Special software (RDSAT) was used to adjust for the chain referral recruitment process used to select the sample for the survey. The software allowed researchers to make adjustments to reflect the make-up of the target population based on information from the networks of the respondents.

Once adjusted, the results were representative of the population, allowing the researchers to compare the results over time. The researchers found a rise in HIV prevalence among men who have sex with men in Beijing over the three years and recommended urgent services for this population to slow the spread of HIV. This study was one of the first to use respondent-driven sampling for basic surveillance – comparing prevalence trends. (41)

### 2.3.6 Convenience sampling

#### **Description**

A convenience sample includes persons in the sample who are easy to reach or convenient to sample. An example might be testing and interviewing all sex workers attending an STI clinic.

#### **Advantages**

Although convenience sampling is not a form of probability sampling, data from convenience samples may be the only information available on populations most at risk for HIV transmission.

#### **Limitations**

In convenience sampling, no rules are used for sample selection. There is no way to estimate error or bias in this sampling scheme.

### 2.3.7 Time–location sampling

#### **Description**

Many of the sampling methods discussed above rely on non-probability methods or tend to overestimate people who are more networked. Time–location sampling methods can avoid these limitations.

This is a probability-based surveillance method that relies on a sampling frame derived from the times when and locations where members of a target population congregate rather than where they live (44).

To conduct time–location sampling:

- List all the venues where the population congregates as the primary sampling unit. For example, the population may meet in certain clubs.
- Randomly select specific times, days and venues.
- The selected venues are visited during the day and time specified.
- Subjects are systematically approached and asked to participate.

This represents a probability-based sample if:

- Every member of the target population has an equal probability of being at the venue at any given time or day.
- Every person selected agrees to participate.
- Everyone gives truthful responses.

Time–location sampling requires good-quality formative research to ensure that no bias is entered into selection of the venue, day and time.

#### **Advantages**

A time–location sample can be derived from key informant interviews which describe where members of the population congregate. Once a full census of these sites has been identified, a random selection of these sites can be chosen for inclusion in the sample. This information will also be useful for prevention activities.

#### **Limitations**

A number of limitations exist.

- The venues selected might not necessarily be frequented by all of the target population.
- It is difficult to estimate the probability of missing someone who does not attend any of the venues.
- Some venues offer little privacy for disclosure of sensitive information.
- The accuracy of self-reported data given in a public setting is questionable.
- Identifying venues that most-at-risk populations frequent could be exposing them to unwanted attention.

Although it is time- and labour-intensive, time–location sampling does have utility for surveillance among groups at higher risk for HIV infection as a result of high-risk behaviour (45).

More guidance on how to conduct a time–location survey can be found at: <http://globalhealthsciences.ucsf.edu/PPHG/assets/docs/tls-res-guide.pdf>.

### Time–location sampling in Harare, Zimbabwe

“Spatial–temporal” sampling, or time–location sampling, was used for a cross-sectional survey in Harare, Zimbabwe of men who frequent beer halls. These men are a proxy group for men who have high-risk sexual behaviour or frequent sex workers. To create the sample, the researchers:

- created a list of all beer halls owned by one operator (which accounted for 75% of the beer halls in Harare),
- created a calendar of 4-hour recruitment events,
- randomly selected the beer halls to fill each calendar event,
- approached every third man who entered the beer hall to assess his eligibility for the survey (eligibility criteria included that the man was not intoxicated).

Interviews were conducted in the privacy of nearby vans and biological samples were taken to test for HIV. Prevention efforts were provided at the same time as the survey. Prevalence of HIV was found to be 30% among men attending beer halls in Harare. Sex while intoxicated, unprotected sex with casual partners, and paying for sex were highly correlated with recent HIV seroconversion. The results of this study suggested that prevention efforts should be focused on patrons of beer halls (46).

### Comparison of time–location sampling and respondent-driven sampling

Time–location sampling and respondent-driven sampling are the most common sampling techniques currently used among populations most-at-risk for HIV. They both have limitations. Sometimes one is preferred over the other (47).

Use respondent-driven sampling when:

- the target population is truly hidden
- it is useful to understand the social connections within the population for programme planning.

Use time–location sampling when:

- the target population is visible
- knowledge of the environment will also assist with programme planning.

### 2.3.8 Sample size

Sample size is an important aspect of the sample design and requires a number of considerations.

- What are the key variables in the study (usually HIV prevalence or behavioural factors)?
- How precise do the estimated results for those variables need to be?
- Does the survey need to be large enough to detect change over time between surveys or differences between key subgroups?

In general, larger sample sizes are better than smaller ones:

- Larger samples provide the ability to distinguish true differences among subgroups of the population or to analyse trends in incidence and risk behaviour. Because most hidden or high-risk populations are themselves heterogeneous, surveillance requires large sample sizes to analyse subgroups.
- Larger samples are needed to make the sample more likely to accurately represent the population (i.e. there is less uncertainty around the estimate).
- It is important to note that a large sample does not necessarily guarantee estimates that are unbiased. It is possible to have a very precise estimate of a biased value.

For more detailed information on calculating sample size, see *Surveillance of populations at high risk for HIV transmission* (48).

### 2.3.9 Sampling strategies for populations most at risk for HIV

#### ***Sex workers***

Sex workers may be working legally or illegally:

- If they are legal and registered, routine screening for sexually transmitted infection may be required. If so, a list of workers may be available from authorities or clinics. In this case, simple random sampling or enumeration may be possible from the list. If no such list is available, time–location sampling may be used. For reasons of comparability, surveillance should be conducted during the same period each year.
- For illegal sex workers who are venue based, time–location sampling at street, brothel, agent or other venues can be used for HIV surveillance. If sex workers are not venue based but are well networked, respondent-driven sampling might be appropriate.

Surveillance in institutional settings such as clinics is less subject to refusal bias, but more subject to participation bias when compared with population-based surveillance. Clinics may overrepresent sex workers at either end of the economic spectrum, depending on whether services are free or not. Some populations may avoid clinics due to legal implications or stigma toward their behaviour.

#### ***Clients of sex workers***

Researchers may use time–location sampling focusing on the locations where persons meet sex workers to provide estimates of behavioural risk and infection prevalence (49). Researchers may also use convenience sampling. In this case, an interviewer approaches clients leaving the room of a female sex worker to invite the client for an interview and to provide an oral fluid sample for rapid testing (15,16). Clients of sex workers are often not well networked so snowball or respondent-driven sampling is not effective in this population.

#### ***Persons who inject drugs***

For out-of-treatment drug users, street- or venue-based sampling or network sampling will be necessary. For in-treatment drug users, convenience samples can be useful. For example, needle exchange programmes can be useful in assessing the changing characteristics of illicit drug markets, even though the data are not representative of the broader population of persons who inject drugs (18). Behavioural surveillance is important to document the effectiveness of prevention programmes in this population (20). Injecting drug users are often well networked in order to purchase drugs so network sampling methods are appropriate.

#### ***Men who have sex with men***

Sampling approaches for HIV surveillance among men who have sex with men again depends on the stigma attached to the behaviour.

Convenience sampling, such as patients at a clinic (50) is often difficult as few countries have clinics devoted to men who have sex with men. In those that do, such clinics are usually limited to large urban centres.

For men who have sex with men, it may be more appropriate to conduct repeated biological and behavioural surveys at venues identified through formative research. If the population is hidden and venues are not available, respondent-driven sampling can be used. Repeating these surveys at consistent sites will be crucial for tracking trends over time.

### Surveillance among men who have sex with men in Bangkok, Thailand

Three cross-sectional surveys were carried out in Bangkok, Thailand between 2003 and 2007. The surveys focused on men who have sex with men using a time–location sampling methodology.

Venues where the men congregated were identified through a mapping exercise followed by a count of the men attending each venue. Venues were then selected for participation in the survey.

HIV prevalence was calculated and regression analyses were used to evaluate differences and trends over time adjusting for time–location cluster sampling. HIV prevalence in 2003, 2005 and 2007 was 17.3%, 28.3% and 30.8%, respectively.

There was a significant increase between 2003 and 2005 but the difference between 2005 and 2007 was not significant. Additional analysis was done by looking at proxy measures of incidence and changes in behaviour over time (*see* section 3.3.5). Based on the prevalence levels and the behavioural data, the authors identified key prevention strategies that would be useful for reversing the epidemic among men who have sex with men in Bangkok (51).

Table 2.2 provides a quick overview of sampling strategies for populations at higher risk for HIV because of high-risk behaviour.

**Table 2.2. Sampling strategies for various populations**

At higher risk	Possible sampling strategies
Sex workers	<ul style="list-style-type: none"> <li>■ Time–location sampling: At hotspots where sex workers sell their trade</li> <li>■ Convenience: List of registered sex workers</li> <li>■ Convenience: List of STI clinics attendees</li> </ul>
Clients of sex workers	<ul style="list-style-type: none"> <li>■ Cluster sampling: survey of proxy populations (truck drivers, fishermen)</li> <li>■ Time–location sampling: hotspots where clients meet sex workers</li> <li>■ Convenience: list of STI clinic attendees reporting sex worker visits</li> </ul>
Persons who inject drugs	<ul style="list-style-type: none"> <li>■ Respondent-driven sampling: starting seeds from an injecting arcade</li> <li>■ Time–location sampling: at street venues where people inject</li> <li>■ Convenience: List of treatment centre attendees</li> </ul>
Men who have sex with men	<ul style="list-style-type: none"> <li>■ Respondent-driven sampling: starting seeds from a club</li> <li>■ Time–location sampling: at venues (bars, parks, etc.)</li> <li>■ Cluster sampling: using surveys of proxy populations (prisoners)</li> </ul>

## Sampling strategies and design tool

A useful tool for determining the most appropriate sampling tool is hosted at [http://globalhealthsciences.ucsf.edu/PPHG/surveillance/CDC-MARPs/sampling\\_selection.htm](http://globalhealthsciences.ucsf.edu/PPHG/surveillance/CDC-MARPs/sampling_selection.htm).

This tool provides sampling strategies based on the population that is being surveyed. It includes most-at-risk populations as well as some populations that are commonly considered proxy populations for surveillance of those most at risk. Figure 2.3 provides a snapshot of the tool and the results for brothel-based sex workers.

**Figure 2.3. Example from the sampling strategies and design tool**


The screenshot shows a web-based interface for the 'Sex Worker' sampling strategy tool. On the left is a vertical navigation menu with the following items: 'Introduction', 'Sampling Methods', 'Sampling Selection Tool', 'Sex Worker', 'MSM', 'Transgender', 'Drug Users', 'Voluntary Migrants', 'Refugees and IDPs', 'STI Clinic Attendees', 'Uniformed Personnel', 'Prisoners', 'Youth', 'Resources', and 'Glossary/Acronyms'. The main content area is titled 'Sex Worker' and is divided into two columns: 'Legal' and 'Illegal/Not registered'. Under 'Legal', there are two options: 'Street-based' and 'Brothel-based'. Under 'Illegal/Not registered', there are two options: 'Agent-based' and 'Venue-based'. Below these options, there are two text boxes: 'No list' and 'List available from brothel owner'. At the bottom of the main content area, there is a text box labeled 'Respondent driven sampling (See Resources)'. At the bottom of the interface, there are two buttons: 'Back' and 'Exit', and a text box labeled 'User Prompt'. The screen number 'Screen: 11Iibii1' is displayed in the bottom right corner.

Developed for Centers for Disease Control and Prevention by Design and Learning Interactive, Inc., [www.designandlearning.com](http://www.designandlearning.com), a Northrop Grumman subcontractor, 2009.

## 3. Conduct surveillance of populations most at risk for HIV

The next part of the process concerns conducting surveys. Populations most at risk for HIV have special requirements. Focus on their special needs and how to avoid legal repercussions and stigma.

**Figure 3.1. Conduct surveillance of most-at-risk populations**



### 3.1 Step 4: Address operational issues

#### 3.1.1 Plan staffing and training

Recruiting and training staff for a surveillance project is a critical and challenging task. The surveillance programme staff should include:

- management and technical personnel
- support staff
- fieldwork staff
- data entry staff
- analysis staff.

Table 3.1 provides a surveillance staff list and job responsibilities. Regular and ongoing training of staff is critical to ensure the success of the surveillance programme. All staff should be aware of all aspects of the programme. Use every opportunity to cross-train staff. For example, give data entry staff field experience in collecting data. They will be better team members.

**Table 3.1 Example responsibilities of surveillance programme staff**

Staff categories	Example responsibilities
Surveillance management and technical personnel	
Surveillance director	<ul style="list-style-type: none"> <li>■ Overall surveillance, statistical and financial direction</li> <li>■ Technical lead for the project</li> </ul>
Surveillance and statistical specialist	<ul style="list-style-type: none"> <li>■ Surveillance methods, statistical survey design, enumeration areas, sampling frames</li> <li>■ Field staff training, supervision, quality control</li> <li>■ Data management, statistical analysis, report writing</li> </ul>
Financial manager	<ul style="list-style-type: none"> <li>■ Surveillance budget accounts</li> <li>■ Spending monitor</li> <li>■ Contracts and acquisitions</li> </ul>
Support staff	<ul style="list-style-type: none"> <li>■ General administrative support for surveillance staff (May include drivers and tracking of fuel and mileage)</li> </ul>
Fieldwork	
Team leader	<ul style="list-style-type: none"> <li>■ Team supervisor</li> <li>■ Meetings with community leader and members</li> </ul>
Interviewer	<ul style="list-style-type: none"> <li>■ Interviewing and obtaining specimens for HIV testing</li> <li>■ Completing questionnaires and edits to questionnaires</li> </ul>
Laboratory technician	<ul style="list-style-type: none"> <li>■ Obtaining specimens for HIV tests if the survey protocol does not allow interviewers to do this</li> <li>■ Performing tests in the laboratory</li> <li>■ Quality control of laboratory data</li> </ul>
Data entry and processing	
Data manager	Manager for: <ul style="list-style-type: none"> <li>■ Data entry, program and database development</li> <li>■ Data editing, quality control</li> </ul>
Data entry personnel	<ul style="list-style-type: none"> <li>■ Questionnaire management</li> <li>■ Final hand-editing</li> <li>■ Data entry from questionnaires into electronic data files</li> </ul>
Analyst	<ul style="list-style-type: none"> <li>■ Supervisor, construction of documentation</li> <li>■ Data analysis</li> <li>■ Interpretation of results for relevant audiences</li> <li>■ Report writing</li> </ul>
Statistician	<ul style="list-style-type: none"> <li>■ Review sample design including sample size calculation</li> <li>■ Consult on analysis and interpretation of results</li> </ul>

### 3.1.2 Develop data collection instruments

When designing a questionnaire, be sure to capture all the variables that are needed. Do not include unnecessary questions in the questionnaire. Consistent questions are important to monitor trends over time, but continuing to ask old questions about treatments that have changed is not useful.

To decide which questions to include in a survey, create a *tabulation plan*. A tabulation plan is a set of mock tables that will be used in the final report. Include exactly which variables will be presented in the analytical report of a survey.

Developing a tabulation plan helps to focus the questions on the necessary data. It also ensures that all of the critical questions needed for the analysis are included in the questionnaire.

When constructing a questionnaire, consider other major data collection efforts within the country. Avoid re-creating questions. Copy word-for-word questions that have been used in previous surveys:

- to allow comparison of results between surveys
- to reduce field-testing of questions.

Make sure the survey questionnaires evolve as the epidemic evolves. Family Health International has developed questionnaires for use among populations most at risk. Consult those examples for ideas (52). (<http://www.fhi.org/en/hivaids/pub/guide/bssguidelines.htm>).

Consider international reporting needs and indicators required for national commitments before finalizing the questionnaire. Also be sure to translate and back-translate the questionnaire to identify any misunderstandings, and pretest the questionnaire in a community outside of the sample.

### 3.1.3 Plan handling of specimens and HIV testing

WHO and UNAIDS have developed guidelines for HIV using testing technologies for surveillance (53). These guidelines consider a country's laboratory infrastructure and surveillance needs, and then suggest methods for selecting, evaluating and implementing HIV testing technologies and strategies. These guidelines provide recommendations for:

- specimen selection
- collection, storage and testing of specimens
- selection and evaluation of appropriate HIV testing strategies
- technologies that meet surveillance objectives
- quality assurance issues.

HIV testing among most-at-risk populations presents unique challenges as well as opportunities. The development of rapid testing technologies allows testing to move from the laboratory to the field. Providing results at the time of the test is very useful when working with populations that are hard to reach, as opportunities for these populations to learn their HIV status might not occur often. Depending on the surveillance design, two strategies have been proposed for most-at-risk populations. See Table 3.2.

**Table 3.2. Testing strategies for most-at-risk populations**

Country/surveillance situation	Testing algorithm
Strategy 1: ■ For unlinked anonymous surveillance – when test results and counselling are not provided to respondents	Use a two-test strategy ■ Both tests need high sensitivity and high negative predictive value. ■ Specimens reactive to both tests are considered HIV-positive.
Strategy 2: ■ For linked surveillance – when test results and counselling are provided to respondents at the time of the survey	Use a three-test strategy ■ Confirmatory results are needed according to the algorithms used for diagnosis. ■ Specimens reactive to three different assays are considered positive.

## 3.2. Step 5: Data management

The reward for careful planning and implementation of surveillance comes with the analysis.

### 3.2.1 Plan the data management system

The structure and content of the database should be specified before data collection begins. Database development involves five steps.

### Procedure: Five steps for developing the survey database

Step	Do this...
1	Determine a system for identification (ID) of records. Make the ID system consistent and comprehensive to uniquely identify each record: <ul style="list-style-type: none"> <li>■ create a unique ID for each participant</li> <li>■ protect personal identity. It should be possible to verify ID numbers and analyse results but not disclose who the participant is. All names, registration numbers or addresses should be deleted.</li> </ul>
2	Design a master file for information across multiple datasets, such as demographic information, which will be included in both behavioural interviews and laboratory data. This allows analysts to link datasets for analysis: <ul style="list-style-type: none"> <li>■ Ensure that the relevant IDs are entered on all forms before data collection.</li> <li>■ Include checks between the different files (for example, birth date).</li> </ul>
3	Design the database. The structure of the database will reflect the various types of data and the timing of data collection. <ul style="list-style-type: none"> <li>■ Include a directory for each dataset collected with separate subdirectories for software programs, datasets and documentation.</li> <li>■ This will ensure that all data files can be traced accurately from data collection through analysis to published paper.</li> </ul>
4	Plan data entry and data analysis. <ul style="list-style-type: none"> <li>■ If the data are collected on paper, ensure that no errors are made during data entry. Develop a reliable data entry system using independent double entry.</li> <li>■ If data are collected electronically, select a reliable transfer method from the machine used for data collection into the project database.</li> <li>■ Regardless of the mode of collection, give data entry staff a list of: <ul style="list-style-type: none"> <li>• acceptable values (that is, valid ranges) for each variable (question item)</li> <li>• fields that are mandatory (for example, subject ID or date of data collection).</li> </ul> </li> </ul>
5	Ensure that the protocol is followed and maintained. Be especially aware of ethical issues when designing the database.

### 3.2.2 Use data management best practices

The purpose of data management is to preserve data integrity between data collection and analysis/reporting (54). Data management best practices are particularly important for HIV surveillance among most-at-risk populations due to the sensitive nature of the data (55,56). Sound data management is an ethical obligation for anyone engaged in data collection.

Family Health International (FHI) has produced guidelines for the management of behavioural data (57). (<http://www.fhi.org/en/HIVAIDS/pub/survreports/firstthingsfirst.htm>)

Begin data verification for quality assurance as data are collected and entered during fieldwork. Use the data entry software program for quality assurance. For example, check all fields to make sure that the values are acceptable and internally consistent.

### 3.2.3 Create a dataset for analysis

Most data management experts recommend that permanent datasets for analyses be created only when data are completely entered, cleaned and frozen; no further changes are allowed.

### 3.2.4 Document the steps used to create the database

In professional data management, all decisions are apparent at every stage, from data collection through analysis.

- Document and explain all the decisions made when cleaning the data.
- Document the steps used to create new variables.
- Document how missing variables are handled.
- Label all variables in each dataset in a manner that conveys information even after the datasets are merged.
- Create codebooks or annotated forms to describe each instrument, map variable names onto the data collected.

### 3.2.5 Maintain confidentiality and security

Surveillance staff must ensure and maintain confidentiality during data entry and processing, and should have been trained in ethical issues important for surveillance.

- All personal identifying information (such as name, identification number, address) should be removed from forms before data entry.
- Store identifying information in locked files or destroy it.
- Do not store computer data files with personal information on computer networks.
- If personal information is stored on a computer, password protect or encrypt it.

Linking behavioural data with HIV status from laboratory data is important for HIV surveillance. However, linking data increases the likelihood of inadvertent disclosure of protected health information.

## 3.3. Step 6: Data analysis

No matter how rigorous the method of analysis, if compromises were made in the surveillance design or data collection, the analysis will be flawed. General guidelines for surveillance analysis can be found elsewhere (58,59). Discussed here are analytical methods for specific issues relating to HIV surveillance among populations most at risk for HIV.

### 3.3.1 Handling non-response

Despite careful data collection, there will be some missing data. Decide how to handle these missing data. For example, for some questions, a *respondent* (the person responding) may not give an answer or items may not apply. Include an explanation of missing data in the database documentation.

There are two ways to deal with missing data:

- Ignore it. That is, analyse the data as they exist and omit them when they are missing. Do this if the team believes that people who do not respond to a particular question are, on average, like those who do respond with respect to other data collected (60). While we might question the wisdom of such an assumption for HIV surveillance, this is the assumption that most statistical analysis packages make.
- If the team agrees that missing data cannot be ignored, consider *imputation methods*. Values are filled in (imputed) for the missing data, based on other data that were supplied by that person (61). Depending on the extent and nature of the missing values, the unadjusted data may produce biased estimates (62).

### 3.3.2 Using weighting

Most surveys require sampling weights to be calculated and applied to adjust the results during analysis and avoid bias. For example, if the sampling strategy has oversampled one subgroup of the population (for example, younger sex workers or brothel-based sex workers), the analyst would need to adjust the general results for the whole population to reflect the true probability of being selected.

Similarly, weighting can adjust for cluster or stratified sampling designs.

Weights can also correct for different response rates. For example, if one subpopulation was more likely to refuse to participate in the survey than other subpopulations, the statistician might adjust the weighting to account for the non-response of this group.

In many surveillance designs and sampling strategies described in this document, it will not be possible to create sampling weights.

- If cluster sampling was used with different probabilities of selection by cluster, it will not be possible to develop weights. Consult a sampling statistician in this case.
- For time–location sampling and respondent-driven sampling methods, sampling weights are used.
  - For respondent-driven sampling, see the tools for respondent-driven sampling analysis at [www.respondentdrivensampling.org](http://www.respondentdrivensampling.org).
  - For time–location sampling, see <http://globalhealthsciences.ucsf.edu/PPHG/assets/docs/tls-res-guide.pdf>.

### 3.3.3 Determining HIV prevalence

To calculate HIV prevalence, divide the total number of people testing HIV-positive by the total number who were tested (33).

Programme managers often require the number of individuals in the population of interest who are infected, even though everyone in the population is not tested during the survey. To make this extrapolation, it is necessary to estimate the size of the population for the denominator. Size estimation has numerous challenges, which are described in the size estimates guidelines (10).

Prevalence data combine new and old infections. HIV prevalence is difficult to interpret in regions where treatment is expanding and thus people who are HIV-positive are living longer. A rising prevalence may be interpreted, as people are living longer because they are receiving ART. Stable or declining HIV prevalence may also be interpreted, as the number of new infections is declining or more infected persons are dying.

In populations most-at-risk for infection, prevalence estimates might also reflect:

- the impact of turnover in populations. For example, new sex workers may enter the industry and older sex workers may leave the industry, or
- sample saturation may occur when the same injecting drug users are tested at a syringe exchange programme over multiple rounds of a survey.

### 3.3.4 Determining HIV prevalence trends

Data on trends provide an important indicator of the trajectory of the epidemic. To be able to accurately compare data and assess trends, the data will need to meet many statistical requirements. These include the following:

- Achieve a sufficient survey sample size to detect a change over time.
- Conduct consistent sample recruitment between surveys. For example:
  - If one survey uses respondent-driven sampling and the next survey uses time–location sampling, the populations in the surveys will be different. Therefore, they are not comparable. However, changing the sampling design to adopt a stronger or more accurate method is encouraged.
  - If non-probability sampling is used, it is not possible to determine whether the sample is representative of the population.
  - If the sample is pulled from a voluntary counselling and testing clinic and there has been a change in the fee structure since the last survey, there could be changes in the population attending the clinic.

Use sites consistently included in the surveillance system over time. If additional sites are added or dropped, it is important to adjust for the fact that prevalence might be systematically different in those sites. Be sure to describe any systematic changes in the final report.

If the samples are comparable over time, conduct tests to see if the trends are statistically significant. The conventional approach to trend analysis is to test the significance of the results through different statistical tests:

- a linear chi-square (63)
- a linear regression (64)
- an exponential regression (65) or
- a categorical (66) model.

Descriptions of these techniques of analyses can be found elsewhere.

### 3.3.5 Determining HIV incidence

When interpreting cross-sectional HIV prevalence data, some people assume that: trends in HIV prevalence reflect trends in transmission, or prevalence is a reasonable surrogate for risk of infection. These assumptions have been shown to be false in some situations (67).

HIV *incidence* (recent infection) is a better measure for understanding the projection of the epidemic and the progress of HIV prevention efforts. However, it is more difficult to measure. Methods include:

- statistical or mathematical modelling (68,69,70,71)
- directly measuring incidence through laboratory assays. These methods are still being developed (72,73,74,75). Since 2006, a series of meetings have been held to discuss progress towards validating a protocol for the use of existing HIV incidence assays and the development of future incidence assays. WHO has established a Technical Incidence Assay Working Group (HIVIWG). Detailed information can be accessed on the WHO web site: ([http://www.who.int/diagnostics\\_laboratory/en/](http://www.who.int/diagnostics_laboratory/en/)).
- proxy measures of incidence. Measuring HIV prevalence trends in a population of newly exposed individuals may be a reasonable proxy for incidence (76). For example, assessing HIV prevalence trends among individuals who have just started injecting drugs or selling sex can be used as a proxy for incidence (77,78).
- longitudinal cohort studies. These studies require large numbers of persons at risk and frequent testing to determine ranges for seroconversion (79,80,81).

### 3.3.6 Dealing with uncertainty

Unless a census is done of the population of interest, the individuals in these surveys will be chosen from larger populations. The individuals might not have the same characteristics as the larger population. Survey results will thus have some level of uncertainty, which should be considered when analysing the results. Describe the uncertainty with the results. Ideally, the statistician on the team would calculate standard errors or confidence intervals around the estimates to indicate the uncertainty of the estimate.

### 3.3.7. Ensuring validity


*Validity* refers to the degree to which the results from the surveillance activity accurately measure the indicator or related public health problem (82). In the context of surveillance of most-at-risk populations, validity is determined primarily by the sample design, sample size, questionnaire construction and accuracy of participants' responses (83).

- Base the sampling strategy or design on a probability sample if possible. If this is not possible, use a strategy that best represents the population.
- Choose the correct sample size to ensure that the data collection is adequate to measure the characteristics of interest. Many populations most at risk for HIV have varied behaviours. Large samples permit analysis of subgroups of the population.
- Reduce bias by constructing a good questionnaire for collecting accurate information from respondents (84).

## 4. Use the results/Evaluate the surveillance plan

The final two steps of the process are using the results and evaluating the surveillance system.

**Figure 4.1. Use the results collected and evaluate the surveillance system**



### 4.1.1 Share the results

Sharing the surveillance results is an ethical imperative. The results of these efforts can have an important impact on the HIV response. Therefore, once the results and recommendations are available, they should be shared in a responsible manner.

Plan to provide the stakeholders involved in the study with the preliminary results. Use their expertise to interpret the data collected.

### 4.1.2 Develop a dissemination plan

Develop a dissemination plan for the stakeholders involved in the planning phase. Consider these questions:

- Who needs to know the results?
- How should the results be presented to each audience?
- When are the results needed by that audience?
- What are the key messages to get across?

Consider sharing the findings in newsletters, which are published faster than peer-reviewed journals. Also consider that the affected community might not have access to journals.

### 4.1.3 Protect the population from stigma or legal action

Ethical questions on how to share the results will vary by the populations affected. Shared surveillance information may cause political pressure in the case of restrictive policies. This may result in harm to the members of risk groups and make it even harder to contact them in future.

It is important that the results are transformed into messages that are clear and not harmful or further stigmatizing to the population.

#### 4.1.4 Use the results to improve programmes

Use the results to inform prevention, care and treatment programmes. The information may have been disseminated widely to a broad group of stakeholders. However, it is critical to take the data to the next step for use by decision-makers.

Make an effort to ensure that programme managers have access to the results to plan appropriate interventions based on the surveillance results. Surveillance results should help improve services for most-at-risk populations and reduce their risk of acquiring HIV infection. Involve the most-at-risk populations not only in the research but also in the development of intervention programmes based on the results.

Analyse data to generate strategic information. Use the information for decision-making and planning. Only in this way can resources devoted to surveillance be justified and used to curb the epidemic in populations most at risk for HIV.

Try to report on a regular basis to communities at risk, partner organizations and national AIDS councils. Support these stakeholders to determine how the results best inform their services. Make this a part of the ongoing activities of surveillance. Surveillance staff should have access to and proactively participate in the HIV response planning process.

Examples of how surveillance data have been used to improve programmes have been highlighted in recent conferences (85).

- In a number of countries in sub-Saharan Africa, surveillance data have been used in simple models (see example below) to estimate where new HIV infections are occurring.
- In Kazakhstan, surveillance data from prison populations have been used to develop an appropriate prevention response for the prison population.
- In Bangladesh, sophisticated subtyping of HIV strains has provided evidence of how HIV has spread between populations and neighbourhoods.

#### Example of use of surveillance data

Surveillance data on most-at-risk populations can be used to improve HIV programmes with “Modes of Transmission” models. Modes of Transmission is a simple spreadsheet model that allows countries to predict where new infections will occur in their country. Using data on HIV prevalence among different subpopulations and most-at-risk populations, a team of epidemiologists from Kenya was able to identify populations where estimates of new infections can be expected. These data were used to help plan interventions in the national strategic plan. This exercise led to a recommendation to increase service provision for men who have sex with men, persons who inject drugs, other drug users and sex workers.

**Figure 4.2: Distribution of new infections in Kenya**



More information on the Modes of Transmission software can be found at [www.unaids.org](http://www.unaids.org).

Source: Kenya National AIDS Control Council. HIV prevention response and modes of transmission analysis. 2009. ([http://www.unaidsrستا.org/files/u1/Kenya\\_MoT\\_Country\\_Synthesis\\_Report\\_22Mar09.pdf](http://www.unaidsrستا.org/files/u1/Kenya_MoT_Country_Synthesis_Report_22Mar09.pdf)).

---

## 4.2. Step 8: Evaluate the system

Any surveillance activity that is implemented must be evaluated regularly to ensure that it is meeting its objectives (86).

More informal routine monitoring is also useful. Reflect on these aspects of the surveillance system:

- Frequency and timeliness of data collection
- Appropriateness of populations under surveillance
- Consistency of sites and groups measured over time
- Coverage/representativeness of the groups for the target populations.

When evaluating the HIV surveillance system, include the following:

- Can the system describe where new infections are occurring?
- Can the system describe how the epidemic is changing?
- Are the data being used to inform programming?
- Are the right data elements being collected?
- Is the choice of methodology appropriate for the situation?
- Does the data collection process adhere to the protocol?
- Has the analysis been conducted with appropriate tools/methods, including weighting, adjustment for design effects, etc.?
- Is data collection restricted to essential data?
- Is there appropriate process documentation?
- Does the activity conducted meet ethical standards?

Look for additional information on surveillance evaluation guidelines on the WHO and UNAIDS web sites in 2011.

# Appendix A: List of additional surveillance resources

Title	Year	Topic
<i>Guidelines on estimating the size of populations most at risk to HIV</i>	2010	Update on methodologies for estimating the size of most-at-risk populations <a href="http://data.unaids.org/pub/Manual/2010/guidelines_popnestimationsize_en.pdf">http://data.unaids.org/pub/Manual/2010/guidelines_popnestimationsize_en.pdf</a>
Evaluation guidelines for HIV second generation surveillance system	Forth-coming 2011	How to assess and evaluate HIV surveillance systems and adapt them to the needs of countries
Ethical issues in HIV surveillance: guidelines	Forth-coming 2011	Ethical principles for conducting HIV surveillance activities
<i>Guidelines for using HIV testing technologies in surveillance: selection, evaluation and implementation: 2009 update</i>	2009	An update in guidance on selecting and utilizing appropriate HIV tests for surveillance purposes <a href="http://www.who.int/hiv/pub/surveillance/hiv_testing_technologies_surveillance_.pdf">http://www.who.int/hiv/pub/surveillance/hiv_testing_technologies_surveillance_.pdf</a>
<i>HIV triangulation resource guide: synthesis of results from multiple data sources for evaluation and decision-making</i>	2009	Guidelines on conducting triangulation, users' manual developed with examples based on experiences in Africa. <a href="http://www.who.int/hiv/pub/surveillance/hiv_triangulation_guide.pdf">http://www.who.int/hiv/pub/surveillance/hiv_triangulation_guide.pdf</a>
<i>The pre-surveillance assessment: guidelines for planning serosurveillance of HIV, prevalence of sexually transmitted infections and the behavioural components of second generation surveillance of HIV</i>	2005	Tools for preparing to implement second generation surveillance including defining and selecting risk groups, sites and feasibility of methods <a href="http://www.who.int/hiv/pub/surveillance/sti/en/index.html">http://www.who.int/hiv/pub/surveillance/sti/en/index.html</a>
<i>Guidelines for measuring HIV prevalence in population-based surveys</i>	2005	Guidelines for national population-based surveys with HIV testing <a href="http://data.unaids.org/pub/Manual/2005/20050101_GS_GuideMeasuringPopulation_en.pdf">http://data.unaids.org/pub/Manual/2005/20050101_GS_GuideMeasuringPopulation_en.pdf</a>
<i>Guidelines for HIV surveillance among tuberculosis patients</i>	2004	How to conduct HIV surveillance among TB patients <a href="http://www.searo.who.int/LinkFiles/Facts_and_Figures_05_GuidelinesHIV_surveillanceTBpatients_final_version.pdf">http://www.searo.who.int/LinkFiles/Facts_and_Figures_05_GuidelinesHIV_surveillanceTBpatients_final_version.pdf</a>
<i>Guidelines for effective use of data from HIV surveillance systems</i>	2004	Guidance on how to analyse, interpret and present surveillance data <a href="http://data.unaids.org/Publications/IRC-pub06/JC1010-UsingData_en.pdf">http://data.unaids.org/Publications/IRC-pub06/JC1010-UsingData_en.pdf</a>

# Appendix B: Glossary and acronyms

<b>AIDS</b>	Late stage of HIV infection, including the development of one or more opportunistic illnesses (illnesses that occur because of low levels of CD4 lymphocytes or immunodeficiency)
<b>Active surveillance</b>	A system in which health authority personnel take the lead in identifying and reporting cases, as opposed to “passive surveillance”
<b>AIDS case surveillance</b>	The identification and reporting of persons meeting the definition of an AIDS case to permit public health authorities to track the disease over time. Also known as “AIDS case reporting”
<b>Anonymous</b>	Having no known name or identity. For example, removing all personal identifying information from a sample that will be tested for HIV, in order to protect the patient’s identity
<b>Antiretroviral drugs</b>	Drugs used to fight infections caused by retroviruses such as HIV
<b>Antiretroviral therapy (ART)</b>	Treatment with drugs that inhibit the ability of HIV to multiply in the body
<b>Behavioural surveillance</b>	Surveys of behaviour that may result in acquiring HIV. This may involve asking a sample of people about risk behaviours such as sexual practices and drug use.
<b>Bias</b>	A systematic error in the collection or interpretation of data
<b>Bridging populations</b>	Persons belonging to high-risk subpopulations who interact with people at lower risk in the general population, making it more likely for the HIV epidemic to shift from high-risk (“concentrated”) populations to the general population
<b>Case reporting</b>	A surveillance system in which persons who are identified as meeting the case definition are reported to public health authorities
<b>Concentrated HIV epidemic</b>	The epidemic state in which HIV has spread to a high level in a defined subpopulation but is not well established in the general population. This epidemic state is characterized by an HIV prevalence that is consistently >5% in at least one defined subpopulation, but <1% in pregnant women in urban areas.
<b>Confidentiality</b>	Protecting information that concerns a study participant or patient from release to those who do not need to have the information
<b>Convenience sampling</b>	The selection of subjects from a population based on accessibility and availability
<b>Demographic information</b>	The “person” characteristics of epidemiology (usually collected with “place” and “time”) – age, sex, race and occupation – used to characterize the populations at risk
<b>Denominator</b>	The population (or population experience, as in person-years, etc.) at risk in the calculation of a proportion or rate. The denominator is the lower portion of a fraction used to calculate a rate or ratio.
<b>Epidemic</b>	The occurrence of a disease (or other health-related event) at a substantially increased level and at a given period of time in relation to a baseline
<b>Epidemiology</b>	The study of the distribution and determinants of health-related states or events in specified populations, and the application of this study to the control of health problems
<b>Generalized HIV epidemic</b>	The epidemic state in which HIV is firmly established in the general population. This epidemic state is characterized by an HIV prevalence that is consistently >1% in pregnant women.
<b>High-risk behaviours</b>	Behaviours that increase the risk of a person contracting a disease
<b>Highly active antiretroviral therapy (HAART)</b>	The use of at least three antiretroviral drugs in combination to suppress viral replication and progression of HIV disease by reducing the viral load to undetectable levels
<b>HIV</b>	Acronym for human immunodeficiency virus

<b>Human immunodeficiency virus (HIV)</b>	A retrovirus that causes AIDS by infecting T-cells of the immune system
<b>HIV counselling and testing (HCT)</b>	A programme that provides both counselling and testing services to communities, allowing persons who are tested to obtain emotional and medical support before and after their HIV tests
<b>Incidence</b>	A measure of the frequency with which an event, such as a new case of illness, occurs in a population over a period of time. The denominator is the population at risk; the numerator is the number of new cases occurring during a given time period.
<b>Informed consent</b>	Informed consent is the provision of adequate information to actively decide to consent to a procedure. It protects the person's freedom of choice and respects their autonomy with regard to decisions affecting their body and health.
<b>Laboratory-based reporting</b>	A surveillance system in which the reports of cases come from clinical laboratories
<b>Linked anonymous HIV testing</b>	A person is tested for HIV, and the specimen is labelled with a code without a name or identifiers that could reveal the person's identity. This method is voluntary and requires obtaining informed consent and making the test results available (with appropriate counselling) to the person tested.
<b>Linked confidential HIV testing</b>	A person agrees to have an HIV test with the assurance that the test result will be kept confidential and only selected health-care providers may be informed. This method is voluntary and requires obtaining informed consent and discussing the test results with the person. Linked confidential testing also allows for the collection of more detailed demographic and risk-behaviour information.
<b>Linking</b>	Refers to whether a tested individual's name or identifying information is associated with their HIV test results
<b>Low-level HIV epidemic</b>	The epidemic state in which HIV has not reached significant levels in any subpopulation, although HIV infection may have existed for many years. This epidemic state is characterized by an HIV prevalence that has not consistently exceeded 5% in any defined subpopulation. This state suggests that networks of risk are rather diffuse or that the virus has only recently been introduced.
<b>MSM</b>	Acronym for men who have sex with men
<b>Natural history of disease</b>	Acronym for men who have sex with men
<b>Odds ratio</b>	A measure of the strength of an association between exposure and disease, calculated by taking the ratio of the odds of disease with the exposure to those odds without exposure. For a rare disease, the odds ratio is a reasonable approximation of the relative risk. Odds ratios are commonly used for case-control studies and result easily from logistic regression analysis.
<b>Passive surveillance</b>	A system in which a health-care provider or worker notifies the health authority of any cases of diseases, as opposed to "active surveillance"
<b>Random sample</b>	A sample derived by selecting individuals such that each individual has the same probability of selection
<b>Proxy population</b>	A population in which a large proportion of the members have a behaviour that puts individuals at increased risk for HIV. Proxy populations are used when it is not possible to specifically sample the most-at-risk population.
<b>Relative risk</b>	A measure of the association between a disease or condition and a factor under study, calculated by dividing the incidence rate among those exposed to the factor by the incidence rate among those not exposed to the factor
<b>Representative sample</b>	A sample whose characteristics correspond to those of the original or reference population
<b>Representativeness</b>	The extent to which the sample resembles the true population
<b>Selection bias</b>	A systematic error in the process of respondent selection for a study or survey
<b>Sensitivity</b>	The proportion of persons with disease who are correctly identified by a screening test or case definition as having the disease
<b>Sentinel case reporting</b>	Reporting cases of a disease from sentinel sites
<b>Sentinel populations</b>	Populations that are subject to sentinel surveillance activities. They may not necessarily be representative of the general population, but they might be the first to be affected by HIV. Examples include persons with sexually transmitted infection or truck drivers.

<b>Sentinel sites</b>	Sites at which sentinel surveillance activities take place, including clinics attended by individuals who may not be representative of the general population but represent groups initially infected or at higher risk for infection than the general population
<b>Sentinel surveillance</b>	A surveillance system that samples populations from facilities or communities for a short period of time on a regular basis
<b>Seroprevalence surveys</b>	Surveys that estimate the HIV prevalence by testing blood for HIV antibody
<b>Serosurveillance</b>	Collecting blood samples for surveillance. Latent, subclinical infections and carrier states can be detected, in addition to clinical cases. This is especially important in the case of HIV and other sexually transmitted infections, which often have a long latent period before symptoms become apparent.
<b>Sexually transmitted infection (STI)</b>	Diseases that are spread by the transfer of organisms from person to person during sexual contact
<b>Simple random sampling</b>	A sampling method that requires the use of a random number table or other method (for instance, computer-based) to generate random numbers that identify the persons to be included in the sample
<b>Surveillance</b>	The systematic collection, analysis, interpretation and dissemination of health data on an ongoing basis, to gain knowledge of the pattern of disease occurrence and potential in a community, in order to control and prevent disease in the community
<b>Systematic sampling</b>	A sampling method that consists of randomly selecting the initial patient who meets the inclusion criteria and then selecting every "nth" (for example, third or fifth) eligible patient thereafter until the predetermined sample size is reached or the survey period is over
<b>Unlinked anonymous testing</b>	In unlinked anonymous testing, a sample of blood originally collected for other purposes is tested for HIV after all information that could identify the source of the blood is eliminated from the sample.
<b>Validity</b>	The degree to which a measurement actually measures or detects what it is supposed to measure

# Appendix C: Links

## **UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance**

Global surveillance of HIV/AIDS and sexually transmitted infections is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, is the coordination and implementation mechanism for UNAIDS and WHO to compile and improve the quality of data needed for informed decision-making and planning at national, regional and global levels. [http://www.unaids.org/en/HIV\\_data/Epidemiology/epiworkinggrp.asp](http://www.unaids.org/en/HIV_data/Epidemiology/epiworkinggrp.asp)

A good selection of key HIV/AIDS surveillance reports and documents include the following:

- *New strategies for HIV/AIDS surveillance in resource-constrained countries* ([http://data.unaids.org/Topics/Epidemiology/Reports/hivsurveillanceconference-addis\\_en.pdf](http://data.unaids.org/Topics/Epidemiology/Reports/hivsurveillanceconference-addis_en.pdf))
- *Ethical issues to be considered in second generation surveillance* ([http://www.who.int/hiv/pub/epidemiology/en/sgs\\_ethical.pdf](http://www.who.int/hiv/pub/epidemiology/en/sgs_ethical.pdf))
- *Guidelines for conducting HIV sentinel serosurveys among pregnant women and other groups* ([http://data.unaids.org/publications/irc-pub06/jc954-anc-serosurveys\\_guidelines\\_en.pdf](http://data.unaids.org/publications/irc-pub06/jc954-anc-serosurveys_guidelines_en.pdf))
- *HIV/AIDS epidemiological surveillance update for the WHO African Region 2002* (<http://www.who.int/hiv/pub/epidemiology/pubafro2003/en/>)
- *Guidelines for second generation HIV surveillance* ([www.who.int/hiv/pub/surveillance](http://www.who.int/hiv/pub/surveillance))

## **Epidemiological fact sheets on HIV/AIDS and sexually transmitted infections**

These fact sheets provide country-specific information on numerous key indicators relevant to HIV.

## **Journal supplements specifically on HIV surveillance**

*AIDS* 2001 Supplement 3

*AIDS* 2005 Supplement 2 on the first global HIV surveillance conference in Addis Ababa, Ethiopia

*Journal of HIV/AIDS Surveillance and Epidemiology* 2010 issue 2 on the second global HIV surveillance conference in Bangkok, Thailand

# References

- 1 Teutsch SM. Considerations in planning a surveillance system. Chapter 2. In: Teutsch SM, Churchill RE, eds. *Principles and practice of public health surveillance*. 2nd edition. New York, Oxford University Press, 2000.
- 2 Fairchild AL et al. Public goods, private data: HIV and the history, ethics, and uses of identifiable public health information. *Public Health Reports*, 2007, 122:7–15.
- 3 WHO/UNAIDS. Ethical issues in HIV surveillance: guidelines. (In press).
- 4 Rhodes SD et al. Exploring Latino men’s HIV risk using community-based participatory research. *American Journal of Health Behavior*, 2007, 31:146–158.
- 5 Council for International Organizations of Medical Sciences. *The international ethical guidelines for biomedical research involving human subjects: guideline 7*. Geneva, WHO, 2002.
- 6 Emanuel EJ et al. Undue inducement in clinical research in developing countries: is it a worry? *Lancet*, 2005, 366:336–340.
- 7 Schenk K, Williamson J. *Ethical approaches to gathering information from children and adolescents in international settings: guidelines and resources*. Washington, DC, Population Council, 2005.
- 8 Baker MG, Fidler DP. Global public health surveillance under new international health regulations. *Emerging Infectious Diseases*, 2006, 12:1058–1065.
- 9 Baral S et al. Elevated risk for HIV infection among men who have sex with men in low- and middle-income countries 2000–2006: a systematic review. *PLoS Medicine*, 2007, 4:e339. doi:10.1371/journal.pmed.0040339.
- 10 *Introduction to sampling most-at-risk populations: sampling strategies and design tool*. Available at: <http://www.theagencyfordesign.com/clients/cdc/marps> (accessed 12 March 2011).
- 11 UNAIDS. *A framework for monitoring and evaluating HIV prevention programmes for most-at-risk populations*. Geneva, UNAIDS, 2008. Available at: [http://data.unaids.org/pub/Manual/2008/jc1519\\_framework\\_for\\_me\\_en.pdf](http://data.unaids.org/pub/Manual/2008/jc1519_framework_for_me_en.pdf) (accessed 28 October 2010).
- 12 UNAIDS/WHO. *Guidelines on estimating the size of populations most at risk to HIV*. Geneva, UNAIDS/WHO, 2010. Available at: [http://data.unaids.org/pub/Manual/2010/guidelines\\_popnestimationsize\\_en.pdf](http://data.unaids.org/pub/Manual/2010/guidelines_popnestimationsize_en.pdf) (accessed 28 October 2010).
- 13 Ghys PD, Jenkins C, Pisani E. HIV surveillance among female sex workers. *AIDS*, 2001, 15 (suppl 3):S33–S40.
- 14 Subramanian T et al. HIV, sexually transmitted infections and sexual behaviour of male clients of female sex workers in Andhra Pradesh, Tamil Nadu and Maharashtra, India. *AIDS*, 2008, 22:S69.
- 15 Vuylsteke BL et al. HIV prevalence and risk behavior among clients of female sex workers in Abidjan, Côte d’Ivoire. *AIDS*, 2003, 17:1691–1694.
- 16 Lowndes CM et al. Management of sexually transmitted diseases and HIV prevention in men at high risk: targeting clients and non-paying sexual partners of female sex workers in Benin. *AIDS*, 2000, 14:2523–2534.
- 17 Des Jarlais DC, Dehne K, Casabona J. HIV surveillance among injecting drug users. *AIDS*, 2001, 15 (suppl 3):s13–s22.

- 18 Topp L et al. Representativeness of injecting drug users who participate in HIV surveillance. *Journal of Acquired Immune Deficiency Syndromes*, 2008, 47:632-638.
- 19 Lau JTF et al. Changes in the prevalence of HIV-related behaviors and perceptions among 1832 injecting drug users in Sichuan, China. *Sexually Transmitted Diseases*, 2008, 35:325-335.
- 20 Lansky A et al. Developing an HIV behavioral surveillance system for injecting drug users: the national HIV behavioral surveillance system. *Public Health Reports*, 2007, 122(S1):48-55.
- 21 Hammett TM et al. Patterns of HIV prevalence and HIV risk behaviors among injection drug users prior to and 24 months following implementation of cross-border HIV prevention interventions in northern Vietnam and southern China. *AIDS Education and Prevention*, 2006, 18:97-115.
- 22 CDC. Trends in HIV/AIDS diagnoses among men who have sex with men—33 states, 2001–2006. *MMWR Morbidity and Mortality Weekly Reports*, 2008, 57:681-686.
- 23 McFarland W, Caceres CF. HIV surveillance among men who have sex with men. *AIDS*, 2001, 15 (suppl 3):S23-S32.
- 24 Family Health International. *Surveillance and assessment reports*. Available at: <http://www.fhi.org/en/HIVAIDS/pub/survreports/index.htm> (accessed 12 March 2011).
- 25 Cloete A et al. Stigma and discrimination experiences of HIV-positive men who have sex with men in Cape Town, South Africa. *AIDS Care*, 2008, 20:1105-1110.
- 26 Smith A et al. Men who have sex with men and HIV/AIDS in sub-Saharan Africa. *Lancet*, 2009, 374:416-422. DOI:10.1016/S0140-6736(09)61118-1.
- 27 Teplin LA et al. HIV and AIDS risk behaviors in juvenile detainees: implications for public health policy. *American Journal of Public Health*, 2003, 93:906-911.
- 28 Dolan K, Kite B, Black E, et al. HIV in prison in low-income and middle-income countries. *Lancet*, 2007, 7:32-41.
- 29 Rotily M et al. Surveillance of HIV infection and related risk behavior in European prisons: a multicentre pilot study. *European Journal of Public Health*, 2001, 11:243-250.
- 30 WHO. *WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification of HIV-related disease in adults and children*. Geneva, WHO, 2007. Available at: <http://www.who.int/hiv/pub/guidelines/hivstaging/en/index.html> (accessed on 12 March 2011).
- 31 Iguchi M et al. Interpreting HIV seroprevalence data from a street-based outreach program. *Journal of Acquired Immune Deficiency Syndromes*, 1994, 7:491-499.
- 32 American Statistical Association. *Survey research methods section*. Available at: <http://www.amstat.org/sections/srms/Proceedings/> accessed on 28 March 2011.
- 33 WHO. *Guidelines for measuring national HIV prevalence in population-based surveys*. Geneva, WHO, 2006. Available at: <http://www.who.int/hiv/pub/surveillance/guidelinesmeasuringpopulation.pdf> (accessed 12 March 2011).
- 34 WHO. *Guidelines for conducting behavioural surveillance*. New Delhi, WHO SEARO, 2001. Available at: [http://www.searo.who.int/LinkFiles/Publications\\_aids-123.pdf](http://www.searo.who.int/LinkFiles/Publications_aids-123.pdf) (accessed on 12 March 2011).
- 35 Ostrow DG, Kessler RC. *Methodological issues in AIDS behavioral research*. New York, Plenum Press, 1993.
- 36 Foreman EK. *Survey sampling principles*. New York, Marcel Dekker, 1991.
- 37 Munro HL et al. Prevalence and determinants of HIV and sexually transmitted infections in a general population-based sample in Mysore district, Karnataka state, southern India. *AIDS*, 2008, 22 (suppl 5):s117-s125.
- 38 Som RK. *Practical sampling techniques*. Boca Raton, FL, CRC Press, 1996.
- 39 Faugier J, Sargeant BA. Sampling hard to reach populations. *Journal of Advanced Nursing*, 2008, 26:790-797.

- 40 Heckathorn DD. Respondent-driven sampling: a new approach to the study of hidden populations. *Social Problems*, 1997, 44:174–199.
- 41 Heckathorn DD. Extensions of respondent-driven sampling: analyzing continuous variables and controlling for differential recruitment. *Sociological Methodology*, 2007, 37:151–207.
- 42 Ma X et al. Trends in prevalence of HIV, syphilis, hepatitis C, hepatitis B, and sexual risk behavior among men who have sex with men: results of 3 consecutive respondent-driven sampling surveys in Beijing, 2004 through 2006. *Journal of Acquired Immune Deficiency Syndromes*, 2007, 45:581–587.
- 43 Goel S, Salganik MJ. Assessing respondent-driven sampling. *Proceedings of the National Academy of Sciences*, 2010, 107:6743–6747.
- 44 Robbins CL et al. Multicity HIV seroprevalence in street youth, Ukraine. *International Journal of STD & AIDS*, 2010, 21:489–496.
- 45 Stueve A et al. Time–space sampling in minority communities: results with young Latino men who have sex with men. *American Journal of Public Health*, 2001, 91:922–926.
- 46 Fritz KE et al. The association between alcohol use, sexual risk behavior, and HIV infection among men attending beerhalls in Harare, Zimbabwe. *AIDS and Behavior*, 2002, 6:221–228.
- 47 Johnson L, Sabin K, Prybylski D. Update for sampling most-at-risk and hidden populations for HIV biological and behavioral surveillance. *Journal of HIV/AIDS Surveillance & Epidemiology*, 2010, 2:2.
- 48 WHO. *Surveillance of populations at high risk for HIV transmission*. Module 6. New Delhi, WHO Regional Office for South-East Asia, 2007. Available at: [http://www.searo.who.int/LinkFiles/Publications\\_Module-6.pdf](http://www.searo.who.int/LinkFiles/Publications_Module-6.pdf) (accessed on 12 March 2011).
- 49 Subramanian T et al. HIV, sexually transmitted infections, and sexual behavior of male clients of female sex workers in Andhra Pradesh, Tamil Nadu and Maharashtra, India: results of a cross-sectional survey. *AIDS*, 2008, 22 (suppl 5):s69–s79.
- 50 Soto RJ et al. Sentinel surveillance of sexually transmitted infections/HIV and risk behaviors in vulnerable populations in 5 Central American countries. *Journal of Acquired Immune Deficiency Syndromes*, 2007, 46:101–111.
- 51 van Griensven F et al. Trends in HIV prevalence, estimated HIV incidence, and risk behavior among men who have sex with men in Bangkok, Thailand, 2003–2007. *Journal of Acquired Immune Deficiency Syndromes*, 2010, 53:234–239. 210.1097/ QAI.1090b1013e3181c1092fc1086.
- 52 Family Health International. Behavioral surveillance surveys: guidelines for repeated behavioral surveys in populations at risk of HIV. Arlington, FHI, 2000. Available at: <http://www.fhi.org/en/hivaids/pub/guide/bssguidelines.htm> (accessed on 6 December 2010).
- 53 UNAIDS/WHO Working Group on Global HIV/AIDS and STD Surveillance. *Guidelines for using HIV testing technologies in surveillance: selection, evaluation and implementation: 2009 update*. Geneva, UNAIDS/WHO, 2009. Available at: [http://www.who.int/hiv/pub/surveillance/hiv\\_testing\\_technologies\\_surveillance\\_.pdf](http://www.who.int/hiv/pub/surveillance/hiv_testing_technologies_surveillance_.pdf) (accessed 6 December 2010).
- 54 Dean AG. Computerizing public health surveillance systems. In: Teutsch SM, Stroup DF, eds. *Principles and practice of public health surveillance*. 2nd edition. New York, Oxford University Press, 2000.
- 55 Payne L. Information sharing and assessment (ISA): can data management reduce risk? *Children & Society*, 2004, 18:383–386.
- 56 Mohania M, Ananthanarayanan R, Gupta A. Some issues in privacy data management. *Data & Knowledge Engineering*, 2007, 63:591–596.
- 57 Family Health International. *First things first: guidelines on management and coding of behavioural surveillance data*. Arlington, FHI, 2006. Available at: <http://www.fhi.org/NR/rdonlyres/em3sbmf74plnwbkg3y64zrajqbktch66cjjvijog3xiyit44muehtnaayszwyq7u6acvbdkrmaxkj/datamanagementguideline1.pdf> (accessed on 6 December 2010).
- 58 Janes GR et al. Descriptive epidemiology: analyzing and interpreting surveillance data. In: Teutsch SM, Stroup DF, eds. *Principles and practice of public health surveillance*. 2nd edition. New York, Oxford University Press, 2000.

- 59 Brookmeyer R, Stroup DF (eds). *Monitoring the health of populations: statistical principles and methods for public health surveillance*. New York, Oxford University Press, 2004.
- 60 Groves RM et al. *Survey nonresponse*. New York, Wiley and Sons, 2002.
- 61 Harrison KM et al. Risk factor redistribution of the national HIV/AIDS surveillance data: an alternative approach. *Public Health Reports*, 2008, 123:618–627.
- 62 Marston M, Jarris K, Slaymaker E. Non-response bias in estimates of HIV prevalence due to the mobility of absentees in national population-based surveys: a study of nine national surveys. *Sexually Transmitted Infections*, 2008, 84:i71–i77.
- 63 Gouws E et al. The epidemiology of HIV infection among young people aged 15–24 years in southern Africa. *AIDS*, 2008, 22 (suppl 4):s5–s16.
- 64 Van der Loeff MFS et al. Sixteen years of HIV surveillance in a West African research clinic reveals divergent epidemic trends of HIV-1 and HIV-2. *International Journal of Epidemiology*, 2006, 35:1322–1328.
- 65 Gouws E, and the International Group on Analysis of Trends in HIV Prevalence and Behaviours in Young People in Countries most Affected by HIV. Trends in HIV prevalence and sexual behaviour among young people aged 15–24 years in countries most affected by HIV. *Sexually Transmitted Infections*, 2010, 86 (Suppl 2):72–83.
- 66 Fleiss JL. The comparison of proportions for many samples. In: *Statistical methods for rates and proportions*. New York, John Wiley; 1973:99–102.
- 67 McFarland W et al. Detection of early HIV infection and estimation of HIV incidence using a sensitive/less sensitive testing strategy at anonymous counseling and testing sites in San Francisco. *Journal of Acquired Immune Deficiency Syndromes*, 1999, 22:484–489.
- 68 Hallett TB et al. Estimating incidence from prevalence in generalised HIV epidemic: methods and validation. *PLoS Medicine*, 2008, 5:e80. doi:10.1371/journal.pmed.0050080.
- 69 Gregson S et al. Demographic approaches to the estimation of incidence of HIV-1 infection among adults from age-specific prevalence data in stable endemic conditions. *AIDS*, 1996, 10:1689–1697.
- 70 Williams B et al. Estimating HIV incidence rates from age prevalence data in epidemic situations. *Statistics in Medicine*, 2001, 20:2003–2016.
- 71 Stover J et al. Estimated HIV trends and program effects in Botswana. *PLoS ONE*, 2008, 3:e3729.
- 72 Parekh BS, McDougal JS. Application of laboratory methods for estimation of HIV-1 incidence. *Indian Journal of Medical Research*, 2005, 121:510–518.
- 73 Janssen RS et al. New testing strategy to detect early HIV-1 infection for use in incidence estimates and for clinical and prevention purposes. *Journal of the American Medical Association*, 1998, 280:42–48.
- 74 Busch MP et al. Beyond detuning: ten years of progress and new challenges in the development and application of assays for cross-sectional HIV incidence estimation, *AIDS*, 2010, 24:2763–2771.
- 75 Guy R et al. Accuracy of serological assays for detection of recent infection with HIV and estimation of population incidence: a systematic review. *Lancet Infectious Diseases*, 2009, 9:747–759.
- 76 Zaba B, Boerma T, White R. Monitoring the AIDS epidemic using HIV prevalence data among young women attending antenatal clinics: prospects and problems. *AIDS*, 2000, 14:1633–1645.
- 77 Garfein RS et al. Prevalence and incidence of hepatitis C virus infection among young adult injection drug users. *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology*, 1998, 18(suppl 1):s11–s19.
- 78 Mahy M et al. A proxy measure for HIV incidence among populations at increased risk to HIV. *Journal of HIV/AIDS Surveillance and Epidemiology*, 2010, 2:8.
- 79 Kingsley LA et al. Temporal trends in human immunodeficiency virus type 1 seroconversion 1984–1989. A report from the Multicenter AIDS Cohort Study (MACS). *American Journal of Epidemiology*, 1991, 134:331–339.

- 80 Des Jarlais DC et al. Continuity and change within an HIV epidemic: injecting drug users in New York City, 1984 through 1992. *Journal of the American Medical Association*, 1994, 271:121–127.
- 81 Wambura M et al. HIV prevalence and incidence in rural Tanzania: results from 10 years of follow-up in an open-cohort study. *Journal of Acquired Immune Deficiency Syndromes*, 2007, 46:616–623.
- 82 United States Population Council. *AIDSQuest: The HIV/AIDS Survey Library Appendix E: selected bibliography of research on validity and reliability of HIV/AIDS-related survey instruments*. Available at: <http://www.popcouncil.org/Horizons/ORToolkit/AIDSQuest/index.html> (accessed on 28 March 2011).
- 83 Boerma JT Sommerfelt AE. Demographic and health surveys (DHS): contributions and limitations. *World Health Statistics Quarterly*, 1993, 46:222–226.
- 84 Seha AM, Klepp KI, Ndeki SS. Scale reliability and construct validity: a pilot study among primary school children in Northern Tanzania. *AIDS Education and Prevention*, 1994, 6:524–534.
- 85 Mahy M, et al. Maximizing the use of surveillance data: innovative examples from around the world. *Journal of HIV/AIDS Surveillance & Epidemiology*, 2010, 2:11.
- 86 UNAIDS. *National AIDS programmes: a guide to monitoring and evaluation*. Geneva, UNAIDS, 2000. Available at: [http://data.unaids.org/publications/IRC-pub05/jc427-mon\\_ev-full\\_en.pdf](http://data.unaids.org/publications/IRC-pub05/jc427-mon_ev-full_en.pdf) (accessed on 12 March 2011).









For more information, contact:  
World Health Organization  
Department of HIV/AIDS  
Avenue Appia 20  
1211 Geneva 27  
Switzerland  
E-mail: [hiv-aids@who.int](mailto:hiv-aids@who.int)  
[www.who.int/hiv](http://www.who.int/hiv)

